

Werk

Label: Article

Jahr: 1983

PURL: https://resolver.sub.uni-goettingen.de/purl?311067255_0019|log24

Kontakt/Contact

[Digizeitschriften e.V.](#)
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

✉ info@digizeitschriften.de

A NOTE ON ARGUESIAN LATTICES

by ALAN DAY*)

(Received May 5, 1981)

§ 0 INTRODUCTION

In [4] Jónsson introduced the Arguesian lattice identity which reflects precisely Desargues' Theorem for projective spaces qua lattices. In this note we "geometrize" Jónsson's equation by proving that one can assume more geometrical facts about the lattice variables in the equational or the (equivalent) implicational form. An interesting consequence of this is the fact that 2-distributive (modular) lattices are Arguesian. This fact reflects the property that one dimensional projective spaces geometrically trivial.

§ 1 PRELIMINARIES

The Arguesian identity and its equivalents have been developed by Jónsson et al in a series of papers (see especially [3], [6] and [7]). An early result, [5], was that Arguesian lattices are modular. Therefore we will assume throughout this paper that all lattices are modular.

Let L be a (modular) lattice. A triangle (or trilateral) in L is an arbitrary triple $a = (a_0, a_1, a_2) \in L^3$. For two triangles, a and b , in L we require certain derived polynomials and statements

1. Definition. For a, b in L

$$(a) p(a, b) = (a_0 \vee b_0) \wedge (a_1 \vee b_1) \wedge (a_2 \vee b_2)$$

$$(b) c_i(a, b) = (a_j \vee a_k) \wedge (b_j \vee b_k), \quad \{i, j, k\} = \{0, 1, 2\}$$

$$(c) CP(a, b) : (a_0 \vee b_0) \wedge (a_1 \vee b_1) \leq a_2 \vee b_2$$

$$(d) AP(a, b) : c_2(a, b) \leq c_0(a, b) \vee c_1(a, b)$$

$CP(a, b)$ is an abbreviation of central perspectivity for the triangles (of points in a projective plane) a and b . $AP(a, b)$ is an abbreviation for the axial perspectivity of these triangles (of points).

*) This research has been supported by the NSERC, Operating Grant A-8190.

2. Definition. A lattice $(L; \vee, \wedge)$ is called:

(a) Arguesian if for any pair of triangles a, b in L , $p(a, b) \leq a_0 \vee \vee (b_0 \wedge (b_1 \vee (c_2 \wedge (c_0 \vee c_1))))$.

(b) Desarguean if for any pair of triangles a, b in L , $CP(a, b)$ implies $AP(a, b)$.

(c) Arguesian (Desarguean) for a given class of triangles if (a) (respectively (b)) holds for that particular class of triangles.

The following theorem is a synopsis of the pertinent results that appear in aforementioned papers.

3. Theorem. ([3], [6], and [7]): Let L be a modular lattice, then the following are equivalent:

(1) L is Arguesian.

(2) L is Desarguean.

(3) L is Desarguean for triangles a, b in L satisfying (A): $a_i \vee p = b_i \vee p = a_i \vee b_i$, $i = 0, 1, 2$.

(4) L is Arguesian for triangles a, b in L satisfying (A): $a_i \vee p = b_i \vee p = a_i \vee b_i$, $i = 0, 1, 2$.

The other notion we need is that of an n -diamond. This notion reflects the geometrical property of $n + 1$ points in general position in a projective space of dimension $\geq n - 1$.

4. Definition. (i) A sequence $d = (d_0, d_1, \dots, d_n)$ in a (modular) lattice L is called an n -diamond if

$$(D_n1) \bigvee_{j \neq i}^{0, n} d_j = v, \quad i \leq n$$

$$(D_n2) d_i \wedge \bigvee_{\substack{k \neq i \\ k \neq j}}^{0, n} d_k = u, \quad i \neq j \leq n.$$

We will call a sequence, d in L , a lower n -diamond (upper n -diamond) if it satisfies (D_n2) (respectively (D_n1)).

(ii) A lattice is called n -distributive if it does not contain an n -diamond.

By Huhn [2] and Hermann & Huhn [1] we know that (modular) n -distributive lattices form an equational class of (modular) lattices and that an n -diamond is either non-trivial ($d_i \neq d_j$ for $i \neq j$) or completely trivial (i.e. $d_0 = d_1 = \dots = d_n$).

The last notion we need is that of an independent set $\{a_1, \dots, a_n\}$ in L .

$$\perp (a_1, \dots, a_n) \quad \text{iff} \quad u = a_i \wedge \bigvee_{j \neq i}^{1, n} a_j \quad \text{for all } i \leq n.$$

§ 2 THE RESULTS

Our goal is to discover what other restrictions one can place on the triangles and still ensure the Desarguean implication. Basically, we are attempting to "regeometrize" the Desarguean implication by removing all "degenerate" cases of variable substitutions. Our first result is an easy reduction to independent triangles.

1. Lemma. *For a modular lattice L , L is Arguesian if and only if L is Desarguean for triangles \mathbf{a} and \mathbf{b} in L satisfying (A) $a_i \vee p = b_i \vee p$ ($= a_i \vee b_i$) $i \leq 2$ and (I) $\perp(a_0, a_1, a_2)$ and $\perp(b_0, b_1, b_2)$.*

Proof: The condition is clearly necessary so let \mathbf{a} and \mathbf{b} be arbitrary centrally perspective triangles in L satisfying (A). By defining

$$x'_i = (x_i \vee x_j) \wedge (x_i \vee x_k)$$

for $x \in \{a, b\}$ and $\{i, j, k\} = \{0, 1, 2\}$ the reader can easily check that \mathbf{a}' and \mathbf{b}' are centrally perspective triangles in L that satisfy (A) and (I). By the condition we infer $\text{AP}(\mathbf{a}', \mathbf{b}')$ and since $c'_i = c_i$ for all i we obtain $\text{AP}(\mathbf{a}, \mathbf{b})$.

Later results will be proven in a similar way as we add more conditions on our triangles. Let us first note however that we may always assume that the triangles \mathbf{a} and \mathbf{b} are *strictly* independent. That is x_i and x_j will be incomparable if $i \neq j$, $x \in \{a, b\}$,

2. Lemma. *Let \mathbf{a} and \mathbf{b} be centrally perspective triangles in a modular lattice L . If either $\{a_0, a_1, a_2\}$ or $\{b_0, b_1, b_2\}$ contains comparable elements, then \mathbf{a} and \mathbf{b} are axially perspective.*

Proof: By symmetry we need only consider the cases $a_0 \leq a_1$, $a_0 \leq a_2$ and $a_0 \leq a_1$. If $a_0 \leq a_1$ then $\text{CP}(\mathbf{a}, \mathbf{b})$ gives us

$$a_0 \leq (a_0 \vee b_0) \wedge (a_1 \vee b_1) \leq a_2 \vee b_2$$

and

$$\begin{aligned} c_0 \vee c_1 &= (a_1 \vee a_2) \wedge (b_1 \vee b_2 \vee c_1) = \\ &= (a_1 \vee a_2) \wedge (b_1 \vee ((b_2 \vee a_0 \vee a_2) \wedge (b_0 \vee b_2))) \geq \\ &\geq (a_1 \vee a_2) \wedge (b_1 \vee ((a_0 \vee b_0) \wedge (a_1 \vee b_1) \wedge (b_0 \vee b_2))) = \\ &= (a_1 \vee a_2) \wedge (a_1 \vee b_1) \wedge (b_1 \vee ((a_0 \vee b_0) \wedge (b_0 \vee b_2))) \geq \\ &\geq a_1 \wedge (b_1 \vee b_0) = \\ &= (a_0 \vee a_1) \wedge (b_0 \vee b_1) = \\ &= c_2. \end{aligned}$$

If $a_2 \leq a_0$ then $\text{CP}(\mathbf{a}, \mathbf{b})$ gives us

$$(a_2 \vee a_1 \vee b_1) \wedge (a_0 \vee b_0) \leq a_2 \vee b_2$$

and

$$\begin{aligned}
c_0 \vee c_1 &= c_0 \vee (a_0 \wedge (b_0 \vee b_2)) = \\
&= (a_0 \wedge (b_0 \vee b_2)) \vee (a_2 \wedge b_0 \vee b_2) \vee c_0 = \\
&= (a_0 \wedge (b_0 \vee b_2)) \vee \{(a_1 \vee a_2) \wedge [b_1 \vee b_2 \vee (a_2 \wedge (b_0 \vee b_2))]\} = \\
&= (a_0 \wedge (b_0 \vee b_2)) \vee \{(a_1 \vee a_2) \wedge [b_1 \vee ((b_2 \vee a_2) \wedge (b_0 \vee b_2))]\} \geq \\
\geq (a_0 \wedge (b_0 \vee b_2)) \vee \{(a_1 \vee a_2) \wedge [b_1 \vee ((a_2 \vee a_1 \vee b_1) \wedge (a_0 \vee b_0) \wedge (b_0 \vee b_2))]\} = \\
&= (a_0 \wedge (b_0 \vee b_2)) \vee \{(a_1 \vee a_2) \wedge [b_1 \vee b_0 \vee (a_0 \wedge (b_0 \vee b_2))]\} = \\
&= [a_1 \vee a_2 \vee (a_0 \wedge (b_0 \vee b_2))] \wedge (b_0 \vee b_1 \vee (a_0 \wedge (b_0 \vee b_2))) \geq \\
&\geq [a_1 \vee ((a_2 \vee b_0 \vee b_2) \wedge a_0)] \wedge (b_0 \vee b_1) \geq \\
&\geq [a_1 \vee [a_0 \vee (b_0 \vee ((a_0 \vee b_0) \wedge (a_1 \vee b_1)))] \wedge (b_0 \vee b_1) = \\
&= [a_1 \vee (a_0 \wedge (a_1 \vee b_0 \vee b_1))] \wedge (b_0 \vee b_1) = \\
&= c_2.
\end{aligned}$$

The case where $a_0 \leq a_2$ is left for the reader.

Now let \mathbf{a} and \mathbf{b} be centrally perspective triangles in L satisfying (A) and (I) define

$$\text{and (i) } s_{\mathbf{a}} = (a_0 \wedge (a_2 \vee b_2)) \vee (a_1 \wedge (a_2 \vee b_2))$$

$$\text{(ii) } s_{\mathbf{b}} = (b_0 \wedge (a_2 \vee b_2)) \vee (b_1 \wedge (a_2 \vee b_2))$$

$$\text{(iii) } a'_i = a_i \vee s_{\mathbf{a}} \text{ and } b'_i = b_i \vee s_{\mathbf{b}}$$

Easy calculations give us that \mathbf{a}' and \mathbf{b}' are centrally perspective triangles satisfying (A) and (I). Moreover $c'_2 = c_2$, $c'_1 = c_1 \vee (c_0 \wedge (a_2 \vee b_2))$ and $c'_0 = c_0 \vee (c_1 \wedge (a_2 \vee b_2))$. Therefore $\text{AP}(\mathbf{a}, \mathbf{b})$ holds if and only if $\text{AP}(\mathbf{a}', \mathbf{b}')$ holds.

The above allows us to add the conditions $a_0 \wedge (a_2 \vee b_2) = a_1 \wedge (a_2 \vee b_2) = 0_{\mathbf{a}}$ and $b_0 \wedge (a_2 \vee b_2) = b_1 \wedge (a_2 \vee b_2) = 0_{\mathbf{b}}$ ($0_{\mathbf{a}} = a_0 \wedge a_1 = a_0 \wedge a_2 = a_1 \wedge a_2$ and similarly for $0_{\mathbf{b}}$) to (A) and (I). We are however more interested in certain consequences of these conditions.

3. Theorem. *A modular lattice, L , is Arguesian if and only if L is Desarguean for triangles \mathbf{a} and \mathbf{b} in L satisfying (A), (I) and (U): $p = (a_i \vee b_i) \wedge (a_j \vee b_j)$, $i \neq j$.*

Proof: Using the notation above the theorem statement we get

$$a'_0 \wedge (a'_2 \vee b'_2) = 0_{\mathbf{a}} \vee s_{\mathbf{a}} \leq p'$$

Therefore $p' = p' \vee (a'_0 \wedge (a'_2 \vee b'_2)) = (p' \vee a'_0) \wedge (a'_2 \vee b'_2)$ since $\text{CP}(\mathbf{a}', \mathbf{b}') = (a'_0 \vee b'_0) \wedge (a'_2 \vee b'_2)$ by (A).

We want one more condition on our triangles. Let \mathbf{a}, \mathbf{b} be triangles in L satisfying (A), (I) and (U) and define

$$\text{(i) } u_{\mathbf{a}} = \bigvee_i^{0,2} (a_i \wedge p), \quad u_{\mathbf{b}} = \bigvee_i^{0,2} (b_i \wedge p),$$

$$\text{(ii) } a'_i = a_i \vee u_{\mathbf{a}}, \quad b'_i = b_i \vee u_{\mathbf{b}}.$$

We have by easy calculations:

- (1) $a'_i \vee b'_i = a_i \vee b_i$
- (2) $p' = p$ and a', b' satisfy (U)
- (3) $a'_i \vee a'_j = a_i \vee a_j \vee (a_k \wedge (a_i \vee b_i)) = a_i \vee a_j \vee (a_k \wedge (a_j \vee b_j))$
- (4) $(a'_i \vee a'_j) \wedge (a'_i \vee a'_k) = [(a_i \vee a_j) \wedge (a_i \vee a_k)] \vee u_a$
- (5) a, b satisfy (I)
- (6) $a'_i \vee p' = a'_i \vee b'_i = b'_i \vee p'$ hence a, b satisfy (A)
- (7) $c'_i = c_i \vee (c_j \wedge (a_k \vee b_k)) = c_i \vee (c_k \wedge (a_j \vee b_j))$
- (8) $AP(a, b)$ iff $AP(a', b')$

Therefore it behooves one to determine what other special properties hold for these triangles.

4. Lemma. (a'_0, a'_1, a'_2, p') and (b'_0, b'_1, b'_2, p') are lower 3-diamonds with bottom element u_a (resp. u_b).

Proof:

$$\begin{aligned} a'_0 \wedge (a'_1 \vee a'_2) &= (a_0 \vee (a_1 \wedge p) \vee (a_2 \wedge p)) \wedge (a_1 \vee a_2 \vee (a_0 \wedge p)) = \\ &= (a_0 \wedge (a_1 \vee a_2)) \vee u_a = \\ &= u_a \end{aligned}$$

$$\begin{aligned} a'_0 \wedge (a'_1 \vee p') &= (a_0 \vee (a_1 \wedge p) \vee (a_2 \wedge p)) \wedge (a_1 \vee p) = \\ &= u_a \vee (a_0 \wedge (a_1 \vee p)) = \\ &= u_a \vee (a_0 \wedge (a_1 \vee b_1) \wedge (a_1 \vee a_0 \vee b_0)) = \\ &= u_a \vee (a_0 \wedge (a_1 \vee b_1)) = \\ &= u_a \vee (a_0 \wedge p) = \\ &= u_a \end{aligned}$$

$$\begin{aligned} p' \wedge (a'_0 \vee a'_1) &= p' \wedge (a_0 \vee a_1 \vee (a_2 \wedge p)) = \\ &= (a_2 \wedge p) \vee [(a_0 \vee a_1) \wedge (a_0 \vee b_0) \wedge (a_1 \vee b_1)] = \\ &= u_a \end{aligned}$$

5. Lemma. For each $\{i, j, k\} = \{0, 1, 2\}$, (a'_i, b'_i, c'_j, c'_k) is a lower 3-diamond.

Proof:

$$\begin{aligned} c'_j \wedge (a'_i \vee b'_i) &= (a_i \vee b_i) \wedge (c_j \vee (c_k \wedge (a_i \vee b_i))) = \\ &= [c_j \wedge (a_i \vee b_i)] \wedge [c_k \wedge (a_i \vee b_i)] = \\ &= c'_k \wedge (a'_i \vee b'_i) \end{aligned}$$

$$\begin{aligned} a'_i \wedge (c'_j \vee c'_k) &= [a_i \vee (a_j \wedge p) \vee (a_k \wedge p)] \wedge [c_j \vee c_k] = \\ &= [a_i \vee (a_j \wedge (a_i \vee b_i)) \vee (a_k \wedge (a_i \vee b_i))] \wedge [c_j \vee c_k] = \\ &= [a_i \vee (b_j \wedge (a_i \vee a_j)) \vee (b_i \wedge (a_i \vee a_k))] \wedge [c_j \vee c_k] = \\ &= (b_j \wedge (a_i \vee a_j)) \vee (b_i \wedge (a_i \vee a_k)) \vee (a_i \wedge (c_j \vee c_k)) \end{aligned}$$

$$\begin{aligned}
&= (b_j \wedge (a_i \vee a_j)) \vee (b_i \wedge (a_i \vee a_k)) \vee (a_i \wedge (b_i \vee b_k)) \vee \\
&\vee (a_i \wedge (b_i \vee b_j)) = \\
&= [c_j \wedge (a_i \vee b_i)] \vee [c_k \wedge (a_i \vee b_i)]
\end{aligned}$$

The other calculations are left to the reader.

6. Theorem. *A modular lattice, L , is Arguesian if and only if any triangles \mathbf{a} , \mathbf{b} in L satisfying:*

$$(A) \quad a_i \vee p = b_i \vee p = a_i \vee b_i$$

$$(I) \quad \perp(a_0, a_1, a_2) \text{ and } \perp(b_0, b_1, b_2)$$

$$(U) \quad p = (a_i \vee b_i) \wedge (a_j \vee b_j), i \neq j$$

(LF) (a_0, a_1, a_2, p) , (b_0, b_1, b_2, p) , (a_i, b_i, c_j, c_k) are lower 3-diamonds are axially perspective. Moreover $c_i \vee c_j = c_i \vee c_k = c_j \vee c_k$.

7. Corollary. *2-distributive (modular) lattices are Arguesian.*

Proof: If L is 2-distributive and \mathbf{a} and \mathbf{b} are triangles in L satisfying (A), (I), (U) and (LF) then the 3-diamond generated by the lower 3-diamond (a_2, b_2, c_0, c_1) must be trivial. This gives

$$[c_0 \wedge (a_2 \vee b_2)] \vee [c_1 \wedge (a_2 \vee b_2)] = q_{02} \wedge q_{12} \wedge a \wedge b$$

where

$$q_{ij} = a_i \vee a_j \vee p, \quad a = a_0 \vee a_1 \vee a_2 \quad \text{and} \quad b = b_0 \vee b_1 \vee b_2.$$

Therefore

$$\begin{aligned}
c_0 \vee c_1 &= c_0 \vee c_1 \vee (q_{01} \wedge q_{02} \wedge a \wedge b) = \\
&= [c_0 \vee c_1 \vee (q_{01} \wedge q_{02})] \wedge a \wedge b = \\
&= [c_0 \vee (q_{02} \wedge (c_1 \vee q_{01}))] \wedge a \wedge b \\
&= (c_0 \vee q_{02}) \wedge a \wedge b = \\
&= a \wedge b \cong \\
&\cong c_2.
\end{aligned}$$

§ 3 DISCUSSION

One would obviously have liked to show that we can assume $0_{\mathbf{a}} = 0_{\mathbf{b}}$ and $a = b$. This however is impossible for as Jónsson (private communication) has observed, 2-dimensional Hall–Dilworth gluings of vector space lattices can produce both Arguesian and non-Arguesian lattices. In the non-Arguesian case one needs $0_{\mathbf{a}} \neq 0_{\mathbf{b}}$.

Andras Huhn has asked if there exists a partial (modular) lattice configuration whose exclusion would characterize Arguesian lattices. The author cannot answer this question at present. Its answer should be connected with the projectivity of

the “correct” version of the Desarguean implication. Our version seems to lack that property.

Finally the author feels that one *should* be able to add $a_i \vee a_j \vee p = a_i \vee a_k \vee p$ to the conditions on the triangles. He (obviously) has no proof of this at this time.

REFERENCES

- [1] C. Hermann and A. Huhn, *Lattices of normal subgroups which are generated by frames*, Colloq. Math. Soc. Janos Bolayi 14 (1974), Lattice Theory, 97—136.
- [2] A. Huhn, *Two notes on n -distributive lattices*, Colloq. Math. Soc. Janos Bolayi 14 (1974), Lattice Theory, 137—147.
- [3] G. Grätzer, B. Jónsson and H. Lakser, *The amalgamation property in equational classes of modular lattices*, Pacific J. Math. 45 (1973), 507—524.
- [4] B. Jónsson, *On the representation of lattices*, Math. Scand. 1 (1953), 193—206.
- [5] B. Jónsson, *Modular lattices and Desargues' theorem*, Math. Scand. 2 (1954), 295—314.
- [6] B. Jónsson, *The class of Arguesian lattices in selfdual*, Alg. Univ. 2 (1972), 396.
- [7] B. Jónsson, and G. S. Monk, *Representations of primary Arguesian lattices*, Pacific J. Math. 30 (1969), 95—139.

A. Day
Dept. of Math., Lakehead University
Thunderbay, Ontario, Postal Code P7B5E1
Canada

