

Werk

Label: Table of literature references

Jahr: 1972

PURL: https://resolver.sub.uni-goettingen.de/purl?31311157X_0097|log69

Kontakt/Contact

Digizeitschriften e.V.
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

✉ info@digizeitschriften.de

Remark 5,2. Theorem 5,2 is a Fredholm type theorem for Fredholm-Stieltjes integral equations (5,6). Let us mention that Eq. (5,8) as well as the equation $\Phi - L\Phi = \Phi^\circ$ are not the adjoint equations to (5,6) in the usual sense. We have not a satisfactory description of the dual space V_n' to V_n , which would make it possible to derive the analytic form of the adjoint operator K^* . Nevertheless conditions for solvability of Eq. (5,6) are obtained in a form which is closely related to the well known Fredholm alternative for Fredholm integral equations of the second kind in L_2 -spaces.

References

- [1] *Kurzweil J.*: Generalized Ordinary Differential Equations and Continuous Dependence on a Parameter, Czech. Math. J., 7 (82), (1957), 418–449.
- [2] *Kurzweil J.*: On Integration by Parts, Czech. Math. J., 8 (83), (1958), 356–359.
- [3] *Schwabik Š.*: Stetige Abhängigkeit von einem Parameter und invariante Mannigfaltigkeiten für verallgemeinerte Differentialgleichungen, Czech. Math. J., 19 (94), (1969), 398–427.
- [4] *Dieudonné J.*: Foundations of Modern Analysis, Academic Press, New-York, London, 1960.
- [5] *Hildebrandt T. H.*: Introduction to the Theory of Integration, Academic Press, New-York, London, 1963.
- [6] *Robertson A. P., Robertson W.*: Topological Vector Spaces, Cambridge University Press, 1964.

Author's address: Žitná 25, Praha 1 (Matematický ústav ČSAV v Praze).