

## Werk

**Label:** Article

**Jahr:** 1973

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?31311157X\\_0098|log28](https://resolver.sub.uni-goettingen.de/purl?31311157X_0098|log28)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

ON THE MINIMUM NUMBER OF VERTICES AND EDGES  
IN A GRAPH WITH A GIVEN NUMBER OF SPANNING TREES

LADISLAV NEBESKÝ, Praha

(Received May 31, 1971)

By a graph we shall mean a finite connected undirected graph without loops and multiple edges (for notions and results of graph theory see, for example, [1] or [2]). If  $p, q$  and  $r$  are integers such that  $1 \leq p \leq q \leq r$  and  $2 \leq q$  then by  $D(p, q, r)$  we shall denote the graph with cyclomatic number 2 and with no separating vertex and such that its two vertices of degree 3 are connected to each other by arcs ([2]) of length  $p, q$  and  $r$ ; the graph  $D(p, q, r)$  has of course  $p + q + r - 1$  vertices,  $p + q + r$  edges and  $pq + qr + pr$  spanning trees.

In the following, by  $x$  we shall denote a positive integer other than 2. By  $\alpha(x)$  we denote the smallest number  $y_1$  such that there is a graph having  $y_1$  vertices and  $x$  spanning trees; by  $\beta(x)$  we denote the smallest number  $y_2$  such that there is a graph having  $y_2$  edges and  $x$  spanning trees. Obviously  $\alpha(x) \leq \beta(x) \leq x$ , for any  $x \geq 3$ . The function  $\alpha$  has been studied by J. SEDLÁČEK [3], who also gave an impulse to the rise of the present paper.

The very simple generalization of one of the procedures used in [3] for the estimate of the function  $\alpha$  leads to the following estimate of the function  $\beta$  which is given by graphs with at least one separating vertex: if  $x_1$  and  $x_2$  are integers and  $x_1, x_2 \geq 3$ , then

$$(1) \quad \beta(x_1 x_2) \leq \beta(x_1) + \beta(x_2).$$

By making use of the graph  $D(1, 2, (x - 2)/3)$  and a graph with no separating edge and with two circuits of length 3 and  $x/3$ , J. Sedláček [3] found an upper estimate of the function  $\alpha$  for almost all  $x \equiv 2, 3 \pmod{3}$ . By using the same graphs it is quite readily possible to find an estimate of the function  $\beta$  for the same values of the argument:

$$(2) \quad \text{if } x \equiv 2 \pmod{3}, \quad x \geq 8, \quad \text{then } \beta(x) \leq (x + 7)/3;$$

$$(3) \quad \text{if } x \equiv 3 \pmod{3}, \quad x \geq 9, \quad \text{then } \beta(x) \leq (x + 9)/3.$$

Estimate (3) of course also follows from estimate (1). Upper estimates of the func-

tion  $\beta$  (and hence also the function  $\alpha$ ) for almost all  $x \equiv 1 \pmod{3}$  are given by the following lemma.

**Lemma.** *It holds that:*

- (4) if  $x \equiv 1 \pmod{30}$ ,  $x \geq 91$ , then  $\beta(x) \leq (x + 269)/30$ ;
- (5) if  $x \equiv 16 \pmod{30}$ ,  $x \geq 106$ , then  $\beta(x) \leq (x + 254)/30$ ;
- (6) if  $x \equiv 4 \pmod{30}$ ,  $x \geq 64$ , then  $\beta(x) \leq (x + 206)/30$ ;
- (7) if  $x \equiv 19 \pmod{30}$ ,  $x \geq 79$ , then  $\beta(x) \leq (x + 221)/30$ ;
- (8) if  $x \equiv 7 \pmod{15}$ ,  $x \geq 37$ , then  $\beta(x) \leq (x + 98)/15$ ;
- (9) if  $x \equiv 10 \pmod{15}$ ,  $x \geq 40$ , then  $\beta(x) \leq (x + 110)/15$ ;
- (10) if  $x \equiv 13 \pmod{15}$ ,  $x \geq 43$ , then  $\beta(x) \leq (x + 92)/15$ .

**Proof.** By  $G_1$  we denote the graph with 10 vertices  $a_1, a_2, a_3, a_4, b_1, b_2, b_3, b_4, b_5, c_0$  and 11 edges  $c_0a_1, a_1a_2, a_2a_3, a_3a_4, a_4c_0, c_0b_1, b_1b_2, b_2b_3, b_3b_4, b_4b_5, b_5c_0$ ;  $G_1$  obviously has 30 spanning trees. By  $G_2$  we denote the graph with 6 vertices  $a_1, a_2, a_3, b_1, b_2, b_3$  and with 8 edges  $a_1a_2, a_2a_3, a_3a_1, b_1b_2, b_2b_3, b_3b_1, a_1b_1, a_3b_3$ ;  $G_2$  obviously has 30 spanning trees. By  $G_3$  we denote the graph with 7 vertices  $a_1, a_2, b_1, b_2, b_3, b_4, c_0$  and 8 edges  $c_0a_1, a_1a_2, a_2c_0, c_0b_1, b_1b_2, b_2b_3, b_3b_4, b_4c_0$ ;  $G_3$  obviously has 15 spanning trees. We now construct graphs  $G_4, \dots, G_{10}$  such that in any one of the graphs  $G_i$ ,  $i = 1, 2, 3$ , we select vertices  $v$  and  $w$ , and then complete the respective graph  $G_i$  by  $j - 1$  vertices and  $j$  edges so that the vertices  $v$  and  $w$  are connected to each other by an arc of length  $j$  of which every inner vertex is different from all vertices of the graph  $G_i$ . We obtain the graph  $G_4, \dots, G_{10}$ , by selecting  $i, v, w$  and  $j$  as follows ( $j$  is, of course, always an integer):

$$\begin{aligned}
 G_4: & i = 1, \quad v = a_2, \quad w = b_1, \quad j = (x - 61)/30 \geq 1; \\
 G_5: & i = 1, \quad v = a_2, \quad w = b_2, \quad j = (x - 76)/30 \geq 1; \\
 G_6: & i = 2, \quad v = a_1, \quad w = b_2, \quad j = (x - 34)/30 \geq 1; \\
 G_7: & i = 2, \quad v = a_1, \quad w = a_2, \quad j = (x - 19)/30 \geq 2; \\
 G_8: & i = 3, \quad v = a_1, \quad w = b_1, \quad j = (x - 22)/15 \geq 1; \\
 G_9: & i = 3, \quad v = a_1, \quad w = a_2, \quad j = (x - 10)/15 \geq 2; \\
 G_{10}: & i = 3, \quad v = a_1, \quad w = b_2, \quad j = (x - 28)/15 \geq 1.
 \end{aligned}$$

There is little difficulty in seeing that the numbers of edges of the graphs  $G_4, \dots, G_{10}$  give successively estimations (4)–(10).

**Theorem 1.** *If  $x = 1$ , then  $\alpha(x) = 1, \beta(x) = 0$ ; if  $x$  is one of the numbers 3, 4, 5, 6, 7, 10, 13, 22, then*

$$(11) \quad \alpha(x) = \beta(x) = x;$$