

## Werk

**Label:** Table of literature references

**Jahr:** 1975

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?31311157X\\_0100|log20](https://resolver.sub.uni-goettingen.de/purl?31311157X_0100|log20)

## Kontakt/Contact

Digizeitschriften e.V.  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

Proposition 2.9 and Lemma 2.1,  $S_{a,b} = S_{e,b} = 1_G$  for all  $a, b \in G$ . The converse is obvious.

(ii)  $\Leftrightarrow$  (iii) The proof is quite similar, using the dual of Lemma 4.7.

#### *References*

- [1] *B. Д. Белоусов*: Основы теории квазигрупп и луп, Москва 1967.
- [2] *T. Kepka*: On one class of quasigroups, Časop. pěst. mat. 97 (1972), 347–356.
- [3] *T. Kepka, P. Němec*: *T-quasigroups*. Part II, Acta Univ. Carolinae, Math. et Phys., 12, 2 (1971), 31–49.
- [4] *S. K. Stein*: On a construction of Hosszú, Publ. Math. Debrecen, 6 (1959), 10–14.

*Author's address:* 186 00 Praha 8, Sokolovská 83 (Matematicko-fyzikální fakulta KU).