

## Werk

**Label:** Article

**Jahr:** 1975

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?31311157X\\_0100|log61](https://resolver.sub.uni-goettingen.de/purl?31311157X_0100|log61)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

ON SOME BOUNDARY VALUE PROBLEMS FOR NONLINEAR  
FOURTH ORDER ORDINARY DIFFERENTIAL EQUATIONS

MICHAL ČVERČKO, Košice

(Received January 25, 1974)

The aim of this paper is to investigate the relationship between the existence of functions satisfying differential inequalities and the existence of a solution to the boundary value problems (1), (2), then (1), (3), and (1), (4), where

$$(1) \quad y^{(4)} = h(x, y, y', y'', y'''),$$

$$(2) \quad \begin{aligned} y(a) &= a_0, \\ y'(a) &= a_1, \\ y''(a) &= a_2, \\ y''(d) &= d_2, \end{aligned}$$

$$(3) \quad \begin{aligned} y''(a) &= a_2, \\ y(d) &= d_0, \\ y'(d) &= d_1, \\ y''(d) &= d_2, \end{aligned}$$

$$(4) \quad \begin{aligned} y''(a) &= a_2, \\ y(b) &= b_0, \\ y'(b) &= b_1, \\ y''(d) &= d_2. \end{aligned}$$

The method of G. A. KLAASEN from his article [1] will be used.

Throughout this paper it is assumed that  $R$  is the set of real numbers,  $I = [a, d]$ ,  $a < b < d$ ,  $a_0, a_1, a_2, b_0, b_1, d_0, d_1, d_2$  are from  $R$ ,  $D = I \times R^4$ ,  $h : D \rightarrow R$  is a continuous function.

**Lemma 1.** *If  $h$  is continuous and bounded, then for any numbers  $a_0, a_1, a_2, d_2$  there exists a solution of the boundary value problem (1), (2).*

Proof. Since the corresponding homogeneous boundary value problem

$$(5) \quad y^{(4)} = 0,$$

$$(6) \quad y(a) = y'(a) = y''(a) = y'''(d) = 0$$

has only the trivial solution, there exists the Green function  $G(x, t)$  of the non-homogeneous BVP (6),  $y^{(4)} = r(x)$  so that the investigated BVP is equivalent to the integro-differential equation

$$y(x) = w(x) + \int_a^d G(x, t) h(t, y(t), y'(t), y''(t), y'''(t)) dt,$$

where  $w : I \rightarrow R$ ,  $w(x) = c_1 x^3 + c_2 x^2 + c_3 x + c_4$  is the solution of the BVP (5), (2).

Let us denote

$$\begin{aligned} K_0 &= \sup \{ |G(x, t)| : (x, t) \in I \times I \}, \\ K_1 &= \sup \{ |G_x(x, t)| : (x, t) \in I \times I \}, \\ K_2 &= \sup \{ |G_{xx}(x, t)| : (x, t) \in I \times I \}, \\ K_3 &= \sup \{ |G_{xxx}(x, t)| : (x, t) \in I \times I \setminus \{(t, t) : t \in I\} \}, \\ K &= \max \{ K_i : i \in \{0, 1, 2, 3\} \}, \\ m &= \sup \{ |h(x, y, z, u, v)| : (x, y, z, u, v) \in I \times R^4 \}, \\ M &= \sup \{ |w^{(s)}(x)| : x \in I, s \in \{0, 1, 2, 3\} \}. \end{aligned}$$

In the Banach space  $B$  of all functions defined on  $I$  which have continuous third derivative with the norm defined by  $\|r\| = \max \{ \max |r^{(s)}(t)| : t \in I \} : s \in \{0, 1, 2, 3\}$ , the set  $S = \{r \in B : \|r\| \leq mK(d - a) + M\}$  is closed and convex. The mapping  $T : S \rightarrow B$  defined by

$$(Tr)(x) = w(x) + \int_a^d G(x, t) h(t, r(t), r'(t), r''(t), r'''(t)) dt$$

is continuous as well as compact and maps  $S$  into itself. It then follows from the Schauder Fixed-Point Theorem that  $T$  has a fixed point in  $S$ . The fixed point is a solution of the stated BVP.

**Remark.** Similar statements hold for the BVP (1), (3) and (1), (4). Applying any one of these statements, we shall refer always to Lemma 1.

**Lemma 2.** Assume that 1.  $h$  is continuous;

2. all solutions of the initial value problems for the equation (1) extend to all  $I$  or extend to the interior  $I^\circ$  of  $I$  and are unbounded in neighbourhoods of the points  $a$  and  $d$ ;
3. functions  $h_n : D \rightarrow R$  are continuous for  $n = 1, 2, \dots$ ;

4. on every compact set  $K \subset D$  the sequence  $(h_n | K)_1^\infty$  uniformly converges to  $h | K$ ;
5.  $y_n$  are solutions of the equations  $y^{(4)} = h_n(x, y, y', y'', y''')$  on  $I$ ;
6. sequences  $(y_n)_1^\infty, (y'_n)_1^\infty, (y''_n)_1^\infty$  are uniformly bounded on  $I$ . Then there exists a solution  $y : I \rightarrow R$  of the equation (1) such that there exists a subsequence  $(y_{n_k})_{k=1}^\infty$  of the sequence  $(y_n)_1^\infty$  with  $y_{n_k}^{(i)} \rightarrow y^{(i)}, i = 0, 1, 2, 3$  and the convergence is uniform on  $I$ .

*Proof.* There exists a number  $M$  such that for all positive integers  $n$  and all  $x \in I$  it is  $|y_n''(x)| \leq M$  (assumption 6). Therefore there exist  $x_n \in I^\circ$  such that

$$|y_n''(x_n)| = \left| \frac{y_n''(d) - y_n''(a)}{d - a} \right| \leq \frac{2M}{d - a}.$$

The sequences  $(x_n), (y_n(x_n)), (y'_n(x_n)), (y''_n(x_n)), (y'''_n(x_n))$  are bounded. Hence there exist subsequences  $(x_{n_k}), (y_{n_k}(x_{n_k})), (y'_{n_k}(x_{n_k})), (y''_{n_k}(x_{n_k}))$  and  $(y'''_{n_k}(x_{n_k}))$  which are all convergent. Let us denote the limits of these subsequences respectively by  $x_0, y_0, y'_0, y''_0, y'''_0$ . For the sake of simplicity we denote the subsequences by  $(x_n), (y_n(x_n)), (y'_n(x_n)), (y''_n(x_n)), (y'''_n(x_n))$ . By applying the standard convergence theorem (see [2], page 15) to the vector differential equation

$$(y, y', y'', y''')' = (y', y'', y''', h(x, y, y', y'', y'''))$$

we get that there exists a subsequence  $((y_{n_k}, y'_{n_k}, y''_{n_k}, y'''_{n_k}))_1^\infty$  of  $((y_n, y'_n, y''_n, y'''_n))_1^\infty$  and a solution  $(y, y', y'', y''')$  of this equation satisfying the initial condition

$$(y, y', y'', y''')(x_0) = (y_0, y'_0, y''_0, y'''_0)$$

and for every compact part of  $I^\circ$

$$(y_{n_k}, y'_{n_k}, y''_{n_k}, y'''_{n_k}) \rightarrow (y, y', y'', y''') \text{ for } k \rightarrow \infty$$

and this convergence is uniform. Since the sequence  $(y_{n_k})$  is uniformly bounded, the function  $y$  is not unbounded, and thus extends to all  $I$  and the convergence is uniform on  $I$ .

**Theorem 1.** Assume that 1.  $h$  is continuous and nonincreasing in the second and the third argument;

2. solutions of (1) extend to  $I$  or to  $I^\circ$  and are unbounded in neighbourhoods of the points  $a$  and  $d$ ;

3. there exists a function  $u : I \rightarrow R$  satisfying the inequality

$$(7) \quad u^{(4)} \geq h(x, u, u', u'', u''');$$

4. there exists a function  $v : I \rightarrow R$  satisfying the inequality

$$(8) \quad v^{(4)} \leq h(x, v, v', v'', v''');$$

5.  $u \leq v, u' \leq v', u'' \leq v''$ ;  
 6.  $u(a) \leq a_0 \leq v(a), u'(a) \leq a_1 \leq v'(a), u''(a) \leq a_2 \leq v''(a), u''(d) \leq d_2 \leq v''(d)$ .

Then there exists a function  $y : I \rightarrow R$  which satisfies the boundary value problem (1), (2) and the inequalities

$$u^{(i)} \leq y^{(i)} \leq v^{(i)}, \quad i = 0, 1, 2.$$

Proof. For every positive integer  $n \geq N_0$ , where

$$N_0 = \max \{ \max \{ |u'''(x)| : x \in I \}, \max \{ |v'''(x)| : x \in I \} \},$$

we define functions  $h_{in} : I \times R^4 \rightarrow R$  as follows:

$$(9) \quad h_{1n}(x, y, y', y'', y''') = \begin{cases} h(x, y, y', y'', n) & \text{for } y''' > n \\ h(x, y, y', y'', y''') & \text{for } |y'''| \leq n \\ h(x, y, y', y'', -n) & \text{for } y''' < -n \end{cases}$$

$$(10) \quad h_{2n}(x, y, y', y'', y''') = \begin{cases} h_{1n}(x, y, y', v'', y''') + \frac{y'' - v''}{1 + y'' - v''} & \text{for } y'' > v'' \\ h_{1n}(x, y, y', y'', y''') & \text{for } u'' \leq y'' \leq v'' \\ h_{1n}(x, y, y', u'', y''') - \frac{u'' - y''}{1 + u'' - y''} & \text{for } y'' < u'' \end{cases}$$

$$(11) \quad h_{3n}(x, y, y', y'', y''') = \begin{cases} h_{2n}(x, y, v', y'', y''') & \text{for } y' > v' \\ h_{2n}(x, y, y', y'', y''') & \text{for } u' \leq y' \leq v' \\ h_{2n}(x, y, u', y'', y''') & \text{for } y' < u' \end{cases}$$

$$(12) \quad h_{4n}(x, y, y', y'', y''') = \begin{cases} h_{3n}(x, v, y', y'', y''') & \text{for } y > v \\ h_{3n}(x, y, y', y'', y''') & \text{for } u \leq y \leq v \\ h_{3n}(x, u, y', y'', y''') & \text{for } y < u \end{cases}$$

The functions  $h_{4n}$  are continuous and bounded. According to Lemma 1 there exist solutions of BVP (2) and

$$(1_n) \quad y_n^{(4)} = h_{4n}(x, y_n, y_n', y_n'', y_n'''), \quad x \in I.$$

Now we shall show that  $u'' \leq y_n'' \leq v''$  for any solution  $y_n$  of (1<sub>n</sub>), (2). Suppose that there exists  $x \in I$  such that  $y_n''(x) > v''(x)$ . Since  $y_n''(a) \leq v''(a), y_n''(d) \leq v''(d)$ , there must exist  $x_0 \in I$  at which the function  $y_n'' - v''$  has a positive relative maximum. Thus

$$(y_n'' - v'')(x_0) > 0, \quad (y_n''' - v''')(x_0) = 0, \quad (y_n^{(4)} - v^{(4)})(x_0) \leq 0.$$

However, according to the assumptions 1 and 4 and the definition of functions  $h_{in}$ ,  $i = 1, 2, 3, 4$  we have

$$\begin{aligned}
& y_n^{(4)}(x_0) - v^{(4)}(x_0) \geq h_{4n}(x_0, y_n(x_0), y_n'(x_0), y_n''(x_0), y_n'''(x_0)) - \\
& \quad - h(x_0, v(x_0), v'(x_0), v''(x_0), v'''(x_0)) \geq \\
& \geq h_{3n}(x_0, v(x_0), y_n'(x_0), y_n''(x_0), y_n'''(x_0)) - h(x_0, v(x_0), v'(x_0), v''(x_0), \\
& v'''(x_0)) \geq h_{2n}(x_0, v(x_0), v'(x_0), y_n''(x_0), y_n'''(x_0)) - h(x_0, v(x_0), v'(x_0), \\
& \quad v''(x_0), v'''(x_0)) \geq h_{1n}(x_0, v(x_0), v''(x_0), v'''(x_0), y_n'''(x_0)) - \\
& \quad - h(x_0, v(x_0), v'(x_0), v''(x_0), v'''(x_0)) + \frac{(y_n''' - v''')(x_0)}{1 + (y_n''' - v''')(x_0)} \geq \\
& \quad \geq h(x_0, v(x_0), v'(x_0), v''(x_0), v'''(x_0)) - \\
& \quad - h(x_0, v(x_0), v'(x_0), v''(x_0), v'''(x_0)) + \frac{(y_n''' - v''')(x_0)}{1 + (y_n''' - v''')(x_0)} > 0.
\end{aligned}$$

This contradiction says that our assumption that there exists  $x \in I$  such that  $y_n'''(x) > v'''(x)$  is false. By a similar argument,  $y_n'''(x) \geq u'''(x)$  on  $I$  can be shown. From  $u^{(i)}(a) \leq y_n^{(i)}(a) \leq v^{(i)}(a)$ ,  $i = 0, 1$  and  $u'' \leq y_n'' \leq v''$  we obtain  $u' \leq y_n' \leq v'$  and  $u \leq y_n \leq v$  on  $I$ . Thus the function  $y_n$  is a solution of the equation

$$(13) \quad y^{(4)} = h_{1n}(x, y, y', y'', y''').$$

Taking into account that  $h_{1n}|K_m \rightrightarrows h|K_m$ , where

$$\begin{aligned}
& K_m = \{(x, y, y', y'', y''') \in D : x \in I, \quad u(x) \leq y \leq v(x), \\
& u'(x) \leq y' \leq v'(x), \quad u''(x) \leq y'' \leq v''(x), \quad |y'''(x)| \leq m\}, \quad m = 1, 2, \dots,
\end{aligned}$$

we see that all conditions of Lemma 2 are satisfied and thus there exists a subsequence  $(y_{n_k})$  and a function  $y : I \rightarrow R$  such that  $y_{n_k} \rightarrow y$  uniformly on  $I$  and

$$y^{(4)}(x) = h(x, y(x), y'(x), y''(x), y'''(x)), \quad x \in I.$$

For all  $k$  we have  $y_{n_k}(a) = a_0$  and therefore  $y_{n_k} \rightarrow y$  yields  $y(a) = a_0$ . In a similar way  $y'(a) = a_1$ ,  $y''(a) = a_2$ ,  $y'''(a) = a_3$  can be shown.

**Theorem 2.** Assume that 1.  $h$  is continuous and nonincreasing in the second and nondecreasing in the third argument;

2. the solutions of (1) extend to  $I$  or to  $I^\circ$  and are unbounded in neighbourhoods of the points  $a$  and  $d$ ;
3. there exists a function  $u : I \rightarrow R$  satisfying (7);
4. there exists a function  $v : I \rightarrow R$  satisfying (8);

5.  $u \leq v, u' \geq v', u'' \leq v''$ ;  
 6.  $u''(a) \leq a_2 \leq v''(a), u(d) \leq d_0 \leq v(d), u'(d) \geq d_1 \geq v'(d), u''(d) \leq d_2 \leq v''(d)$ .

Then there exists a function  $y : I \rightarrow R$  which satisfies the boundary value problem (1), (3) and the inequalities

$$u \leq y \leq v, \quad u' \geq y' \geq v', \quad u'' \leq y'' \leq v''.$$

**Proof.** Similarly as in the proof of Theorem 1, let us define the function  $h_{1n}$  by (9),  $h_{2n}$  by (10),  $h_{4n}$  by (12), and  $h_{3n}$  as follows:

$$h_{3n}(x, y, y', y'', y''') = \begin{cases} h_{2n}(x, y, u', y'', y''') & \text{for } y' > u', \\ h_{2n}(x, y, y', y'', y''') & \text{for } v' \leq y' \leq u', \\ h_{2n}(x, y, v', y'', y''') & \text{for } y' < v'. \end{cases}$$

Then the functions  $h_{4n}$  are continuous and bounded. According to Lemma 1 there exist solutions  $y_n$  of BVB (3) and (1<sub>n</sub>).

By the method used in the proof of the preceding theorem,  $u'' \leq y_n'' \leq v''$  for any solution  $y_n$  can be shown. From these inequalities as well as from the inequalities

$$u'(d) \geq y_n'(d) \geq v'(d)$$

we obtain

$$u' \geq y_n' \geq v'.$$

Now the last inequality and the inequality

$$u(d) \leq y_n(d) \leq v(d)$$

yields the inequality

$$u \leq y_n \leq v.$$

Thus  $y_n$  is a solution of the equation (13).

Taking into account that  $h_{1n} | K_m \Rightarrow h | K_m$ , where

$$K_m = \{(x, y, y', y'', y''') \in D : x \in I, \quad u(x) \leq y \leq v(x), \\ u'(x) \geq y' \geq v'(x), \quad u''(x) \leq y'' \leq v''(x), \quad |y'''(x)| \leq m\}, \quad m = 1, 2, \dots,$$

we see that all conditions of Lemma 2 are satisfied and thus there exists a subsequence  $(y_{n_k})$  and a function  $y : I \rightarrow R$  such that  $y_{n_k} \rightarrow y$  uniformly on  $I$  and

$$y^{(4)}(x) = h(x, y(x), y'(x), y''(x), y'''(x)), \quad x \in I.$$

For all  $k$  we have  $y_{n_k}(d) = d_0$  and therefore  $y_{n_k} \rightarrow y$  yields  $y(d) = d_0$ . In a similar way  $y'(d) = d_1, y''(d) = d_2, y''(a) = a_2$  can be shown.

**Theorem 3.** Suppose that 1. the function  $h$  is continuous, nonincreasing in the second argument and nondecreasing in the third argument for each  $x \in [a, b]$  as well as nonincreasing for each  $x \in [b, d]$ ;

2. the solutions of initial value problems for (1) extend to  $I$  or to its interior  $I^\circ$  and are unbounded in neighbourhoods of the points  $a$  and  $d$ ;
3. there exist functions  $u \in C^{(4)}(I)$ ,  $v \in C^{(4)}(I)$  satisfying (7) and (8), respectively.
4.  $u \leq v$ ,  $x \in [a, b] \Rightarrow v'(x) \leq u'(x)$ ,  $x \in [b, d] \Rightarrow u'(x) \leq v'(x)$ ,  $u'' \leq v''$ ;
5.  $u''(a) \leq a_2 \leq v''(a)$ ,  $u(b) = b_0 = v(b)$ ,  $u'(b) = b_1 = v'(b)$ ,  $u''(d) \leq d_2 \leq v''(d)$ .

Then there exists a function  $y : I \rightarrow R$  which satisfies (1), (4) and

$$u \leq y \leq v, x \in [a, b] \Rightarrow v'(x) \leq y'(x) \leq u'(x), x \in [b, d] \Rightarrow u'(x) \leq y'(x) \leq v'(x), \\ u'' \leq y'' \leq v''.$$

Proof. Let us denote  $N_0 = \max \{ \max \{ |u'''(x)| : x \in I \}, \max \{ |v'''(x)| : x \in I \} \}$  and for all  $n \geq N_0$  define the functions  $h_{1n}$  by (9),  $h_{2n}$  by (10) and  $h_{3n}$ ,  $h_{4n}$  as follows:

$$h_{3n}(x, y, y', y'', y''') = \begin{cases} h_{2n}(x, y, v', y'', y''') & \text{for } a \leq x \leq b, \quad y' < v' \text{ and} \\ & b \leq x \leq d, \quad y' > v', \\ h_{2n}(x, y, y', y'', y''') & \text{elsewhere,} \\ h_{2n}(x, y, u', y'', y''') & \text{for } a \leq x \leq b, \quad y' > u' \text{ and} \\ & b \leq x \leq d, \quad y' < u', \end{cases}$$

$$h_{4n}(x, y, y', y'', y''') = \begin{cases} h_{3n}(x, v, y', y'', y''') & \text{for } y > v, \\ h_{3n}(x, y, y', y'', y''') & \text{for } u \leq y \leq v, \\ h_{3n}(x, u, y', y'', y''') & \text{for } y < u. \end{cases}$$

Every function  $h_{4n}$  is continuous and bounded and therefore (see Lemma 1) there exists a  $y_n$  which satisfies (1<sub>n</sub>), (4).

We will now show that  $u''(b) \leq y_n''(b) \leq v''(b)$ . Suppose that  $y_n''(b) < u''(b)$ . Since  $y_n''(a) \geq u''(a)$ ,  $y_n''(d) \geq u''(d)$ , there exists a subinterval containing  $b$  on which  $y_n''(x) - u''(x) < 0$  and in that subinterval there exists an  $x_0$  at which  $y_n'' - u''$  has negative relative minimum (and either  $x_0 < b$  and  $y_n'(x_0) > u'(x_0)$  or  $x_0 \geq b$  and  $y_n'(x_0) \leq u'(x_0)$ ). And thus we have

$$y_n'''(x_0) = u'''(x_0), \quad y_n^{(4)}(x_0) \geq u^{(4)}(x_0).$$

However,

$$\begin{aligned} & y_n^{(4)}(x_0) - u^{(4)}(x_0) \leq h_{4n}(x_0, y_n(x_0), y_n'(x_0), y_n''(x_0), y_n'''(x_0)) - \\ & \quad - h(x_0, u(x_0), u'(x_0), u''(x_0), u'''(x_0)) \leq \\ & \leq h_{3n}(x_0, u(x_0), y_n'(x_0), y_n''(x_0), y_n'''(x_0)) - h(x_0, u(x_0), u'(x_0), u''(x_0), u'''(x_0)) \leq \\ & \leq h_{2n}(x_0, u(x_0), u'(x_0), y_n''(x_0), y_n'''(x_0)) - h(x_0, u(x_0), u'(x_0), u''(x_0), u'''(x_0)) \leq \\ & \leq h_{1n}(x_0, u(x_0), u'(x_0), u''(x_0), y_n'''(x_0)) - h(x_0, u(x_0), u'(x_0), u''(x_0), \\ & \quad u'''(x_0)) - \frac{u''(x_0) - y_n''(x_0)}{1 + u''(x_0) - y_n''(x_0)} = h(x_0, u(x_0), u'(x_0), u''(x_0), u'''(x_0)) - \\ & \quad - h(x_0, u(x_0), u'(x_0), u''(x_0), u'''(x_0)) - \frac{u''(x_0) - y_n''(x_0)}{1 + u''(x_0) - y_n''(x_0)} < 0 \end{aligned}$$