

## Werk

**Label:** Article

**Jahr:** 1977

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?31311157X\\_0102|log58](https://resolver.sub.uni-goettingen.de/purl?31311157X_0102|log58)

## Kontakt/Contact

Digizeitschriften e.V.  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

## REMARK ON THE THEOREM OF EGOROFF

WŁADYSŁAW WILCZYŃSKI, ŁÓDŹ

(Received October 29, 1975)

I. VRKOČ in [1] has proved the following theorem: *there exists a real function  $f$  defined and measurable on  $[0, 1]$  such that there does not exist a countable family  $\{A_n\}$  of sets fulfilling  $\bigcup_n A_n = [0, 1]$  such that the restricted function  $f|_{A_n}$  is continuous for every  $n$ .* The theorem of Vrkoč is a refinement of the well known theorem of Lusin. In this short note we shall prove the theorem which can be considered as a similar refinement of the theorem of Egoroff.

Before stating the theorem we shall prove the following lemma:

**Lemma.** *Let  $\{n(k, i)\}$  be a double sequence of natural numbers, which for every  $k$  is increasing with respect to the variable  $i$ . There exists an increasing sequence  $\{n(i)\}$  of natural numbers such that for every  $k$  and for every  $i \geq k$*

$$n(i) > n(k, i).$$

**Proof.** Put  $n(i) = 1 + \max(n(1, i), n(2, i), \dots, n(i, i))$  for every natural  $i$ . It is easy to see that the sequence  $\{n(i)\}$  fulfills all required conditions.

**Theorem.** *For every set  $A$  of the power of continuum there exists a sequence of real functions  $\{f_n\}$  defined on  $A$  such that  $f_n(x)$  tends to zero for every  $x \in A$  and there does not exist a countable family  $\{A_k\}$  of sets fulfilling  $\bigcup_k A_k = A$  such that the restricted sequence  $\{f_n|_{A_k}\}$  is uniformly convergent for every  $k$ .*

**Proof.** Let  $N$  be a set of all increasing sequences of natural numbers. Of course,  $N$  is a set of the power of continuum. Let  $\Phi: A \rightarrow_{\text{onto}} N$  be a one-to-one correspondence.

For  $x \in A$  let us put  $f_{n(1)}(x) = 1^{-1}$ ,  $f_{n(2)}(x) = 2^{-1}$ , ...,  $f_{n(i)}(x) = i^{-1}$ , ... and  $f_j(x) = 0$  for remaining natural  $j$ , where  $\{n(1), n(2), \dots, n(i), \dots\} = \Phi(x)$ .

So we have defined a sequence of real functions  $\{f_n\}$  and it is easy to verify that  $f_n(x) \rightarrow 0$  for every  $x \in A$ .

Suppose that there exists a sequence  $\{A_k\}$  of sets such that  $\bigcup_k A_k = A$  and  $\{f_n|_{A_k}\}$  tends uniformly to 0 for every  $k$ .