

Werk

Label: Article

Jahr: 1981

PURL: https://resolver.sub.uni-goettingen.de/purl?31311157X_0106|log102

Kontakt/Contact

[Digizeitschriften e.V.](#)
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

✉ info@digizeitschriften.de

SOME REMARKS ON DOMATIC NUMBERS OF GRAPHS

BOHDAN ZELINKA, Liberec

(Received July 26, 1979)

E. J. Cockayne and S. T. Hedetniemi in the papers [1] and [2] define the domatic number of an undirected graph. Here we shall present some results concerning this concept. We shall investigate finite undirected graphs without loops and multiple edges.

A dominating set in a graph G is a subset D of the vertex set $V(G)$ of G with the property that each vertex of $V(G) - D$ is adjacent to at least one vertex of D . A partition of $V(G)$ into dominating sets is called a *domatic partition of G* . The maximal number of classes of a domatic partition of a graph G is called *the domatic number of G* and is denoted by $d(G)$.

In [2] it is suggested to relate the domatic number of a graph G to the connectivity of this graph. In this paper we shall prove some results concerning this topic.

The vertex (or edge) connectivity degree of a graph G is the minimal cardinality of a subset of the vertex set (or the edge set, respectively) of G with the property that by deleting this set from G a disconnected graph is obtained. (To delete a subset of the vertex set of G means to delete all vertices of this set and all edges which are incident to these vertices. To delete a subset of the edge set of G means to delete only all edges of this set.) The vertex connectivity degree of G will be denoted by $\omega(G)$, its edge connectivity degree by $\sigma(G)$.

Theorem 1. *Let p and q be non-negative integers, $p < q$. Then there exists a graph G such that $\omega(G) = p$, $d(G) = q$.*

Proof. Take two copies G' , G'' of the complete graph K_q with q vertices. If $p = 0$, then G is the graph whose connected components are G' and G'' . If $p \neq 0$, we choose pairwise distinct vertices u_1, \dots, u_p in G' and v_1, \dots, v_p in G'' and identify u_i with v_i for each $i = 1, \dots, p$. In the following we shall denote the vertex obtained by identifying u_i with v_i by w_i for $i = 1, \dots, p$. The remaining vertices of G' (or G'') will be denoted by u_{p+1}, \dots, u_q (or v_{p+1}, \dots, v_q , respectively). In the case $p = 0$ we denote the vertices of G' by u_1, \dots, u_q and the vertices of G'' by v_1, \dots, v_q . If we delete the set $\{w_1, \dots, w_p\}$ from G , we obtain a disconnected graph. As each of the vertices w_1, \dots, w_p is adjacent to all the other vertices of G , after deleting less than p vertices

the graph G remains connected; therefore $\omega(G) = p$. Let $D_i = \{w_i\}$ for $i = 1, \dots, p$ and $D_i = \{u_i, v_i\}$ for $i = p + 1, \dots, q$. Evidently $\{D_1, \dots, D_q\}$ is a domatic partition of G and $d(G) \geq q$. In [1] it was proved that $d(G) \leq \delta(G) + 1$, where $\delta(G)$ is the minimal degree of a vertex of G . Here evidently $\delta(G) = q - 1$, hence $d(G) = q$.

Theorem 2. *Let p and q be non-negative integers, $p < q$. Then there exists a graph G such that $\sigma(G) = p$, $d(G) = q$.*

Proof. We take again two copies G' and G'' of K_q . Let the vertices of G' (or G'') be u_1, \dots, u_q (or v_1, \dots, v_q , respectively). If $p = 0$, the graph G is the same as in the proof of Theorem 1. If $p \neq 0$, we join u_i with v_i by an edge for each $i = 1, \dots, p$. Evidently $\sigma(G) = p$, where G is the graph thus obtained. Taking $D_i = \{u_i, v_i\}$ for $i = 1, \dots, q$ we obtain a domatic partition $\{D_1, \dots, D_q\}$ and, as $\delta(G) = q - 1$, we have $d(G) = q$.

Theorem 3. *Let h be a positive integer. Then there exists a graph G such that*

$$\omega(G) - d(G) = \sigma(G) - d(G) = h.$$

Proof. Let $n = 2h + 4$ and consider the complete graph K_n . As n is even, there exists a linear factor F of K_n . Let the edges of F be e_1, \dots, e_{h+2} , let u_i, v_i be the end vertices of the edge e_i for $i = 1, \dots, h + 2$. Let G be the graph obtained from K_n by deleting all edges of F . Evidently each subset of $V(G)$ which induces a disconnected subgraph of G is of the form $\{u_i, v_i\}$ for some i . Therefore $\omega(G) = n - 2 = 2h + 2$. It is easy to prove that also $\sigma(G) = n - 2 = 2h + 2$. No vertex of G is adjacent to all the other vertices, therefore each dominating set of G has at least two vertices. This implies $d(G) \leq n/2$. Putting $D_i = \{u_i, v_i\}$ for $i = 1, \dots, h + 2$ we obtain a domatic partition of G and hence $d(G) = n/2 = h + 2$. We have

$$\omega(G) - d(G) = \sigma(G) - d(G) = h.$$

The graph from the proof of Theorem 3 also has the property that $d(G) = \frac{1}{2} \delta(G) + 1$. We express a conjecture.

Conjecture. *For each graph G we have*

$$d(G) \geq \frac{1}{2} \delta(G) + 1.$$

At the end we turn to another problem suggested in [2] – to characterize the uniquely domatic graphs.

A graph G is called *uniquely domatic*, if there exists exactly one domatic partition of G with $d(G)$ classes.

We shall characterize the uniquely domatic graphs whose domatic number is 2. First we prove a lemma.