

Werk

Label: Article

Jahr: 1976

PURL: https://resolver.sub.uni-goettingen.de/purl?316342866_0017|log27

Kontakt/Contact

[Digizeitschriften e.V.](#)
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

✉ info@digizeitschriften.de

A NOTE ON CLOSED CATEGORIES

Jan PAVELKA, Praha

Abstract: For an adjoint situation
 $\mathcal{V}_0(A \otimes B, C) \approx \mathcal{V}_0(A, [BC])$
 in a category \mathcal{V}_0 , the paper gives a description in terms
 of the left adjoint \otimes of those closed category structures
 in the sense of Eilenberg-Kelly on \mathcal{V}_0 that have
 $[-, -]$ for the internal hom-functor. It turns out that \otimes
 need not really be (even up to an isomorphism) associative.

Key-Words: Adjoint situation, closed category, internal
 hom-functor, natural associativity.

AMS: 18D15

Ref. Ž.: 2.726.14

Introduction. Although the concept of a closed category
 is the minimal one of the enrichments of category theory
 treated in [1], it already provides enough framework for
 some interesting applications (the study of \mathcal{V} -categories,
 \mathcal{V} -functors, etc.). Of course, it facilitates the calculus
 considerably if the internal hom-functor

$$[-, -] : \mathcal{V}_0^* \times \mathcal{V}_0 \longrightarrow \mathcal{V}_0$$

has a left adjoint

$$\otimes : \mathcal{V}_0 \times \mathcal{V}_0 \longrightarrow \mathcal{V}_0$$

Nevertheless, once an adjoint to $[-, -]$ is considered it
 is always required to be (up to a specified natural isomor-

phism) associative.

Since there exist closed categories in which the internal hom-functor has a non-associative adjoint (an example will be given in Section 2), we can ask what it is on the side of \otimes that exactly corresponds to an extension of $[-,-]$ to a closed category structure on \mathcal{V}_0 .

To settle this question we first analyze, on similar lines as in [1], Chapter II, §§ 3, 4, the interaction between properties of \otimes and those of $[-,-]$ induced by the adjunction. This time, however, we shall emphasize whatever independence there is between individual couples of corresponding data or axioms and we shall go as far as possible without normalization of the couple $\langle \mathcal{V}_0, [-,-] \rangle$. As for the statements 1.2, 2.2, 3.2, and 3.3 of Section 1, this results in a certain restriction on the proof techniques available and the proofs are, consequently, longer than those in [1]. Because, on the other hand, their complexity is due only to complexity of the calculus involved and they are based on a very simple idea, we shall illustrate the idea by carrying out one of the proofs in question and omit all others. The proofs of 1.1, 2.1, and 3.1 will be also omitted - the reader can be referred to [1], Chapter II, Lemma 3.1.

Convention: The identity morphism of an object A will often be also denoted by A . We denote by \bar{f} the inverse of an isomorphism f .

We shall constantly refer to diagrams MC1 - MC4 on p. 472 and to diagrams CC1 - CC4 on p. 429 in [1]. When we say, for instance, that some transformations a and ℓ satisfy MC1

it means that every diagram of the sort labelled on p.472 by MCl commutes.

1. Relations between data and axioms. Throughout this section we shall deal with the following basic situation: we assume given bifunctors

$$\otimes : \mathcal{V}_0 \times \mathcal{V}_0 \rightarrow \mathcal{V}_0 \quad \text{and} \quad [-, -] : \mathcal{V}_0^* \times \mathcal{V}_0 \rightarrow \mathcal{V}_0$$

together with a natural isomorphism

$$\pi_{ABC} : \mathcal{V}_0(A \otimes B, C) \approx \mathcal{V}_0(A, [BC]).$$

We shall also use the alternative description of π by its unit: $\epsilon_{AB} = \pi_{A, B, A \otimes B}(A \otimes B) : A \rightarrow [B, A \otimes B]$

natural in A and dinatural in B, and

counit: $\epsilon_{AB} = \pi_{[AB], A, B}([AB]) : [AB] \otimes A \rightarrow B$
 natural in B and dinatural in A.

1.1. Given a transformation

$$(1.1) \quad a_{ABC} : (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C)$$

natural in A, B, C, the formula

$$(1.2) \quad I_{BC}^A = \pi_{[BC] \otimes [AB], A, B} \cdot \pi_{[BC][AB][AC]} \{ \epsilon_{BC} \cdot ([BC] \otimes \epsilon_{AB}) \cdot a_{[BC][AB]A} \}$$

defines a transformation

$$(1.3) \quad I_{BC}^A : [BC] \rightarrow [[AB][AC]]$$

natural in B, C and dinatural in A.

Conversely, given (1.3), the formula

$$(1.4) \quad a_{ABC} = \overline{\pi}_{A, B, [C, A \otimes (B \otimes C)]}$$

$$\begin{aligned} \cdot \pi_{A \otimes B, C, A \otimes (B \otimes C)} \{ [\vartheta_{BC} [C, A \otimes (B \otimes C)]] \cdot \\ \cdot \vartheta_{B \otimes C, A \otimes (B \otimes C)}^C \cdot \vartheta_{A, B \otimes C} \} \end{aligned}$$

defines a transformation (1.1).

Moreover, the procedures (1.2) and (1.4) are mutually inverse and thus establish a 1-1 correspondence between (1.1) and (1.3).

(We shall speak about π -corresponding couples of transformations $\langle a, L \rangle$.)

1.2. Let $\langle a, L \rangle$ be a π -corresponding couple. Then a satisfies MC3 iff L satisfies CC3.

Proof of $CC3 \implies MC3$ (a shortened version): We have to show that under the assumption CC3 the equality

$$(1.5) \quad (A \otimes a_{BCD}) \cdot a_{A(B \otimes C)D} \cdot (a_{ABC} \otimes D) = a_{AB(C \otimes D)} \cdot a_{(A \otimes B)CD}$$

holds for all $A, B, C, D \in \text{obj } \mathcal{V}_0$. Since π is an isomorphism we can as well prove (1.5) with

$$\pi_{A, B, [C[DE]]} \cdot \pi_{A \otimes E, C, [DE]} \cdot \pi_{(A \otimes B) \otimes C, D, E},$$

where $E = A \otimes (B \otimes (C \otimes D))$, applied to both sides. Now

$$\begin{aligned} \pi \pi \pi \{ (A \otimes a_{BCD}) \cdot a_{A(B \otimes C)D} \cdot (a_{ABC} \otimes D) \} = \\ = \pi \pi \{ [D, (A \otimes a) \cdot a \cdot (a \otimes D)] \cdot \vartheta_{(A \otimes B) \otimes C, D} \} = \\ = \pi \pi \{ [D, (A \otimes a) \cdot a] \cdot [D, a \otimes D] \cdot \vartheta \} \end{aligned}$$

which by the naturality of ϑ equals

$$\begin{aligned} & \pi \pi \{ [D, (A \otimes a) \cdot a] \cdot \vartheta_{A \otimes (B \otimes C), D} \cdot a_{ABC} \} = \\ & = [B[C[D, (A \otimes a) \cdot a]]] \cdot [B[C, \vartheta_{A \otimes (B \otimes C), D}]] \cdot \\ & \cdot [B[C, a]] \cdot [B, \vartheta_{A \otimes B, C}] \cdot \vartheta_{AB} \cdot \end{aligned}$$

We apply (1.4) for a_{ABC} and obtain

$$\begin{aligned} & [B[C[D, (A \otimes a) \cdot a]]] \cdot [B[C, \vartheta]] \cdot \\ & \cdot [\vartheta_{BC} [C, A \otimes (B \otimes C)]] \cdot L_{B \otimes C, A \otimes (B \otimes C)}^C \cdot \vartheta_{A, B \otimes C} \cdot \end{aligned}$$

By the naturality of ϑ , the naturality of L (applied three times), and (1.4) for $a_{A(B \otimes C)D}$, the last line can be rewritten as

$$\begin{aligned} & [\vartheta_{BC} [D, A \otimes (B \otimes (C \otimes D))]] \cdot L_{B \otimes C, [D, A \otimes ((B \otimes C) \otimes D)]}^C \cdot \\ & \cdot [B \otimes C [D, A \otimes a]] \cdot [\vartheta_{B \otimes C, D} [D, A \otimes ((B \otimes C) \otimes D)]] \cdot \\ & \cdot L_{(B \otimes C) \otimes D, A \otimes ((B \otimes C) \otimes D)}^D \cdot \vartheta_{A, (B \otimes C) \otimes D} \cdot \end{aligned}$$

Next we use the naturality of L , dinaturality of ϑ , again the naturality of L (three times), then (1.4) for a_{BCD} applied in the first variable of $[-, -]$, and the dinaturality of L , and obtain

$$\begin{aligned} & [\vartheta_{B(C \otimes D), 1}] \cdot [1, [\vartheta_{CD}, 1]] \cdot [L_{C \otimes D, B \otimes (C \otimes D)}^D, 1] \cdot \\ & \cdot L_{[D, B \otimes (C \otimes D)], [D, A \otimes (B \otimes (C \otimes D))]}^{[D, C \otimes D]} \cdot L_{A \otimes (B \otimes C), A \otimes (B \otimes (C \otimes D))}^D \cdot \\ & \cdot \vartheta_{A, B \otimes (C \otimes D)} \cdot \end{aligned}$$

By CC3, this equals

$$[\vartheta, 1] \cdot [1, [\vartheta_{CD}, 1]] \cdot [1, L_{C \otimes D, A \otimes (B \otimes (C \otimes D))}^D] \cdot$$

$$\cdot L_{B \otimes (C \otimes D), A \otimes (B \otimes (C \otimes D))}^{C \otimes D} \cdot \vartheta$$

Using three times the fact that $[-, -]$ is a functor and by (1.4) for ${}^a_{AB}(C \otimes D)$ we obtain

$$\begin{aligned} & [B [\vartheta_{CD, 1}]] \cdot [B, L^D] \cdot [B [C \otimes D, {}^a_{AB}(C \otimes D)]] \cdot \\ & \cdot [B, \vartheta_{A \otimes B, C \otimes D}] \cdot \vartheta_{AB} \end{aligned}$$

which, by the naturality of L applied in the second variable of $[-, -]$, the description of π via ϑ , and by (1.4) for ${}^a_{(A \otimes B)CD}$, yields

$$\begin{aligned} & \pi \{ [C [D, {}^a_{AB}(C \otimes D) \cdot {}^a_{(A \otimes B)CD}]] \cdot [C, \vartheta_{(A \otimes B) \otimes C, D}] \cdot \\ & \cdot \vartheta_{A \otimes B, C} \} = \pi \pi \pi \{ {}^a_{AB}(C \otimes D) \cdot {}^a_{(A \otimes B)CD} \} . \end{aligned}$$

2.1. Let $I \in \text{obj } \mathcal{V}_0$. Then the formulas

$$(2.1) \quad i_A = \pi_{AIA}(r_A) = [I, r_A] \cdot \vartheta_{AI}$$

$$(2.2) \quad r_A = \overline{\pi}_{AIA}(i_A) = \vartheta_{IA} \cdot (i_A \otimes I)$$

establish a 1-1 correspondence between natural transformations

$$(2.3) \quad r_A: A \otimes I \rightarrow A$$

and

$$(2.4) \quad i_A: A \rightarrow [IA]$$

Moreover, r is an isomorphism iff i is.

2.2. Given π -corresponding couples $\langle a, L \rangle$ and $\langle r, i \rangle$. Then a, r satisfy MC4 iff L, i satisfy CC4.

3.1. Let $I \in \text{obj } \mathcal{V}_0$. Then the formulas

$$(3.1) \quad j_A = \pi_{IAA}(l_A) = [A, l_A] \cdot \vartheta_{IA}$$

$$(3.2) \quad l_A = \bar{\pi}_{IAA}(j_A) = e_{AA} \cdot (j_A \otimes A)$$

establish a 1-1 correspondence between

$$(3.3) \quad \text{natural transformations } l_A: I \otimes A \rightarrow A$$

and

$$(3.4) \quad \text{dinatural transformations } j_A: I \rightarrow [AA]$$

3.2. Given π -corresponding couples $\langle a, L \rangle$ and $\langle l, j \rangle$. Then a, l satisfy MC1 iff L, j satisfy CC1.

3.3. Given π -corresponding couples $\langle a, L \rangle, \langle r, i \rangle$, and $\langle l, j \rangle$. Then a, l, r satisfy MC2 iff L, i, j satisfy CC2.

3.4. Given π -corresponding couples $\langle r, i \rangle$ and $\langle l, j \rangle$. Then $r_I = l_I$ iff $i_I = j_I$.

4.1. Given a transformation (3.4) put for any $\xi: A \rightarrow B$ in \mathcal{V}_0

$$(4.1) \quad \tau_{AB}(\xi) = [A, \xi] \cdot j_A$$

We obtain a natural transformation

$$\tau_{AB}: \mathcal{V}_0(A, B) \rightarrow \mathcal{V}_0(I, [AB]).$$

4.2. Let $\langle l, j \rangle$ be a π -corresponding couple and

let τ be defined by (4.1). Then

a) if ℓ is an isomorphism, so is τ and its inverse is determined by

$$(4.2) \quad \bar{\tau}_{AB}(\eta) = e_{AB} \cdot (\eta \otimes A) \cdot \bar{\ell}_A$$

where $\eta : I \rightarrow [AB]$ in \mathcal{V}_0 .

b) If τ is an isomorphism, so is ℓ and we have

$$(4.3) \quad \bar{\ell}_A = \bar{\tau}_{A, I \otimes A}(\vartheta_{IA})$$

Proof: All the verifications are straightforward except perhaps that of 4.2 b. Assume τ is an isomorphism and put

$$(4.4) \quad \hat{\ell}_A = \bar{\tau}_{A, I \otimes A}(\vartheta_{IA}).$$

We show that $\hat{\ell}_A$ is inverse to ℓ_A . For every $A \in \mathcal{V}_0$ we have

$$\ell_A \cdot \hat{\ell}_A = \{ \mathcal{V}_0(A, \ell_A) \cdot \bar{\tau}_{A, I \otimes A} \} \vartheta_{IA} =$$

$$\text{(by the naturality of } \bar{\tau} \text{)} = \{ \bar{\tau}_{AA} \cdot \mathcal{V}_0(I, [A, \ell_A]) \} \vartheta_{IA} =$$

$$= \bar{\tau}_{AA} \{ [A, \ell_A] \cdot \vartheta_{IA} \}$$

$$\text{(by (3.1))} = \bar{\tau}_{AA} \{ j_A \} =$$

$$\text{(by (4.1))} = 1_A.$$

To complete the proof it now suffices to show that each $\hat{\ell}_A$ is an epimorphism. Suppose that

$$A \xrightarrow{\hat{\ell}_A} I \otimes A \xrightarrow[\varphi_1]{\varphi_0} B$$

commutes. Then

$$\pi_{IAB} \varphi_0 = [A, \varphi_0] \cdot \vartheta_{IA} = [A, \varphi_0] \cdot [A, \hat{\ell}_A] \cdot j_A =$$

$$= [A, \varphi_1] \cdot [A, \hat{\ell}_A] \cdot j_A = [A, \varphi_1] \cdot \vartheta_{IA} = \pi_{IAB} \varphi_1$$

whence $\varphi_0 = \varphi_1$.

2. The comparison theorem.

Proposition. In the basic situation of Section 1, the following statements are equivalent:

- (T) \otimes can be extended to a structure on \mathcal{V}_0 whose definition is obtained from the concept of a monoidal category as defined in [1] by weakening the associativity of \otimes to α_{ABC} being just morphisms in \mathcal{V}_0 natural in A, B, C,
 (H) $[-, -]$ can be extended to a closed category structure on \mathcal{V}_0 as defined in [1].

Moreover, the structures on \mathcal{V}_0 mentioned in (T) and (H), respectively, determine each other (up to some freedom we have when defining the basic functor V in the transition from (T) to (H)) uniquely.

Proof: a) (H) \rightarrow (T). Given a closed category structure $\langle V, [-, -], I, L, i, j \rangle$ on \mathcal{V}_0 , use (1.4), (2.2), and (3.2) to obtain transformations α, r, ℓ satisfying MC1 - MC4; r is an isomorphism. By Propositions 2.4 and 2.7 of Chapter I in [1],

$$Vi_{[AB]} = \tau_{AB},$$

where τ is defined by (4.1), holds for any A, B \in obj \mathcal{V}_0 and

$$i_I = j_I.$$

Hence ℓ is an isomorphism and $r_I = \ell_I$ (MC5 in [1]).

b) (T) \mapsto (H). Given $\langle \otimes, I, a, r, \ell \rangle$ such that a, r, ℓ satisfy MC1 - MC5 and r, ℓ are isomorphisms, use (1.2), (2.1), (3.1) and (4.1) to obtain transformations L, i, j, τ such that L, i, j satisfy CC1 - CC4, i, τ are isomorphisms and $i_I = j_I$. It remains to normalize the couple $\langle \mathcal{V}_0, [-, -] \rangle$ (cf. [11, p. 491]). To this end, we prove the following

Lemma. Given i, j such that $i_I = j_I$ and τ is an isomorphism. For any $C \in \text{obj } \mathcal{V}_0$,

$$\begin{cases} \text{if } C = [AB] \text{ put } \mathcal{V}_C = \mathcal{V}_0(AB), \quad \iota_C = \tau_{AB} \\ \text{otherwise put } \mathcal{V}_C = \mathcal{V}_0(I, C), \quad \iota_C = 1_{\mathcal{V}_0(I, C)}. \end{cases}$$

For any $f: C \rightarrow D$ in \mathcal{V}_0 define a mapping $Vf: \mathcal{V}_C \rightarrow \mathcal{V}_D$ by

$$Vf = \tau_D \cdot \mathcal{V}_0(I, f) \cdot \iota_C$$

We obtain a functor $V: \mathcal{V}_0 \rightarrow \text{Set}$ and a natural isomorphism

$$\iota: V \approx \mathcal{V}_0(I, -) \text{ such that}$$

$$(CC0) \quad (i) \quad V \cdot [-, -] = \mathcal{V}_0(-, -)$$

(ii) $V i_{[AB]} = \tau_{AB}$ holds for any $A, B \in \text{obj } \mathcal{V}_0$, in particular,

$$(CC5) \quad V i_{[AA]}(1_A) = j_A$$

Proof: (i) is clearly true on objects. Next, for any

$$A' \xrightarrow{f} A \xrightarrow{\xi} B \xrightarrow{g} B' \text{ in } \mathcal{V}_0$$

$$\begin{aligned} \tau_{A'B'} \{V[f, g] \xi\} &= \{ \tau_{A'B'} \cdot \tau_{A'B'} \cdot \mathcal{V}_0(I, [fg]) \cdot \\ &\cdot \tau_{AB} \} \xi = \{ \mathcal{V}_0(I, [fg]) \cdot \tau_{AB} \} \xi = [fg] \cdot [A \xi] \cdot j_A = \\ &= [f, g \xi] \cdot j_A = [A', g \xi] \cdot [f, A] \cdot j_A = [A', g \xi] \cdot [A', f] \cdot j_{A'} = \\ &= [A', g \xi f] \cdot j_{A'} = \tau_{A'B'}(g \xi f) = \tau_{A'B'} \{ \mathcal{V}_0(f, g) \xi \}, \end{aligned}$$

hence $V[f, g] \xi = \mathcal{V}_0(f, g) \xi$.

(ii) For any $A \xrightarrow{f} B$ in \mathcal{V}_0 we have

$$\begin{aligned} V_{I[AB]} \xi &= \{ \tau_{I, [AB]} \cdot \mathcal{V}_0(I, i_{[AB]}) \cdot \tau_{AB} \} \xi = \\ &= \tau_{I[AB]} \{ i_{[AB]} \cdot [A \xi] \cdot j_A \} = \tau_{I[AB]} \{ [I[A \xi]] \cdot i_{[AA]} \cdot j_A \} = \\ &= \tau_{I[AB]} \{ [I[A \xi]] \cdot [I, j_A] \cdot i_I \} = \tau_{I[AB]} \{ [I[A \xi]] \cdot \\ &\cdot [I, j_A] \cdot j_I \} = \tau_{I[AB]} \{ \tau_{I[AB]} \{ \tau_{AB} \} \} \xi = \tau_{AB} \xi . \end{aligned}$$

Example. To satisfy ourselves that there exist closed categories in which the internal hom-functor has a non-associative adjoint, let us turn to the following special case.

Consider a partially ordered set (i.e. a small thin skeletal category) $\langle P, \leq \rangle$. A closed category structure on $\langle P, \leq \rangle$ boils down to a couple $\langle [-, -], I \rangle$, where $I \in P$ and $[-, -] : P \times P \rightarrow P$ is an operation order reversing in the first and order preserving in the second variable, such that

$$[y, z] \leq [[xy] [xz]]$$

$$x = [Ix]$$

$$I \leq [xx]$$

$$x \leq y \text{ iff } I \leq [x, y]$$

hold for any $x, y, z \in P$.

Now take the closed category structure $\langle [-, -], 3 \rangle$ on the category



where $[-, -]$ is defined by Table 1. Table 2 shows the value of its adjoint \otimes .

3	3	3	3	3
2	3	3	3	2
1	3	3	2	1
0	3	1	1	0
	0	1	2	3

Table 1

3	0	1	2	3
2	0	0	1	2
1	0	0	1	1
0	0	0	0	0
	0	1	2	3

Table 2

Observe that $(2 \otimes 2) \otimes 2 = 1 \otimes 2 = 0 < 1 = 2 \otimes 1 =$
 $= 2 \otimes (2 \otimes 2).$

R e f e r e n c e s

- [1] EILENBERG S. and KELLY G.M.: Closed categories, in Proceedings of the Conference on Categorical Algebra, La Jolla 1965, Springer-Verlag 1966, 421-562.
- [2] KELLY G.M.: Tensor products in categories, J. Algebra 2 (1965), 15-37.
- [3] MacLANE S.: Natural associativity and commutativity, Rice University Studies 49(1963), No 4, 28-46.
- [4] FULTER A.: Extending tensor products to structures of closed categories, Comment. Math. Univ. Carolinae 13(1972), 599-616.
- [5] SCHIPPER de W.J.: Symmetric closed categories, Mathematical Centre Tracts, Mathematisch Centrum, Amsterdam 1975.

Matematicko-fyzikální fakulta
 Karlova universita
 Sokolovská 83, 18600 Praha 8
 Československo

(Oblatum 26.2. 1976)