

Werk

Label: Article

Jahr: 1976

PURL: https://resolver.sub.uni-goettingen.de/purl?316342866_0017|log73

Kontakt/Contact

[Digizeitschriften e.V.](#)
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

✉ info@digizeitschriften.de

DECOMPOSITION OF SPHERES IN HILBERT SPACES

P. ERDÖS, D. PREISS, Budapest - Praha

Abstract: A simple construction of a graph with \aleph_2 vertices and with the chromatic number \aleph_1 whose every subgraph spanned by \aleph_1 vertices has chromatic number $\leq \aleph_0$ is given.

Key word: Chromatic number of a graph.

AMS: 05C15

Ref. Ž.: 8.83

Assume the generalized continuum hypothesis. Consider the unit sphere of the Hilbert space of $\aleph_{\alpha+2}$ dimensions. We join two of its points by an edge if their distance is greater than $\frac{3}{2}$. Since $\frac{3}{2} < \sqrt{3}$ the chromatic number of this graph is by the following theorem $\aleph_{\alpha+1}$ (a graph is called m -chromatic if one can color its vertices by m colors so that two vertices which get the same color are not joined, but one cannot do this with fewer than m colors). On the other hand every subgraph spanned by $\aleph_{\alpha+1}$ vertices has again by the following theorem chromatic number $\leq \aleph_{\alpha}$. A different construction of such graphs is given in [1].

This note was written at the Durham symposium on the relations between infinite-dimensional and finite-dimensional convexity (1975).

Theorem. Let $\aleph_0 \leq n < m$ be cardinal numbers. Then (i) - (iii) are equivalent and imply (iv), moreover, under generalized continuum hypothesis they are equivalent to (iv).

(i) For every $c > \sqrt{2}$ the unit sphere in a Hilbert space of m dimensions can be written as a union of n sets with diameter $< c$.

(ii) There is a number $c \in (\sqrt{2}, \sqrt{3})$ such that the unit sphere in $\mathcal{L}_2(m)$ can be written as a union of n sets with diameter $< c$.

(iii) There is a family \mathcal{C} of subsets of m such that $\text{card}(\mathcal{C}) \leq n$ and \mathcal{C} separates points of m (i.e. for $\alpha, \beta \in m, \alpha \neq \beta$ there is a set $C \in \mathcal{C}$ with $\text{card}(C \cap \{\alpha, \beta\}) = 1$).

(iv) $m \leq 2^n$

Proof. The implications (i) \implies (ii) and (iii) \implies (iv) are obvious. (ii) \implies (iii): Let $\{A_\sigma; \sigma \in n\}$ be sets in $\mathcal{L}_2(m)$ with diameter $< \sqrt{3}$ covering the unit sphere in $\mathcal{L}_2(m)$. For $\alpha, \beta \in m, \alpha \neq \beta$ put

$$\begin{aligned} x_{\alpha, \beta}(\gamma) &= 1/\sqrt{2} \text{ for } \gamma = \alpha \\ &= -1/\sqrt{2} \text{ for } \gamma = \beta \\ &= 0 \text{ otherwise.} \end{aligned}$$

Put $C_\sigma = \{\alpha \in m; \text{there exists } \beta \in m, \beta \neq \alpha \text{ such that } x_{\alpha, \beta} \in A_\sigma\}$.

If $\alpha, \beta \in m, \alpha \neq \beta$ then there is a σ such that $x_{\alpha, \beta} \in A_\sigma$. Consequently, $\alpha \in C_\sigma$ and $\beta \notin C_\sigma$ since $\|x_{\alpha, \beta} - x_{\beta, \gamma}\| \geq \sqrt{3}$ for any γ . Therefore the family $\{C_\sigma; \sigma \in n\}$ separates points in m .

(iii) \implies (i): Let $0 < \varepsilon < \frac{1}{2}$. Let \mathcal{A} be a family of subsets of m separating points of m . We may and will suppose that \mathcal{A} is closed under complements and finite intersections. Let \mathcal{B} be the system of all pairs of finite sequences $\{(A_1, \dots, A_p), (r_1, \dots, r_p)\}$ where $A_1, \dots, A_p \in \mathcal{A}$ are nonempty and disjoint and r_1, \dots, r_p are rational numbers that $1 > \sum_{i=1}^p r_i^2 > (1 - \varepsilon)^2$. For $\sigma \in \mathcal{B}$, $\sigma = \{(A_1, \dots, A_p), (r_1, \dots, r_p)\}$ put $C_\sigma = \{x \in \mathcal{L}_2(m); \|x\| = 1 \text{ and there are } \alpha_i \in A_i \text{ such that } \sum_{i=1}^p (x(\alpha_i) - r_i)^2 < \varepsilon^2\}$. First prove that the family $\{C_\sigma; \sigma \in \mathcal{B}\}$ covers the unit sphere in $\mathcal{L}_2(m)$. If $x \in \mathcal{L}_2(m)$, $\|x\| = 1$ find $\alpha_1, \dots, \alpha_p$ such that $\|y - x\| < \varepsilon$ where $y(\alpha_i) = x(\alpha_i)$ and $y(\alpha) = 0$ for all other α . Since \mathcal{A} is closed under complements and finite intersections, we can find disjoint sets $A_i \in \mathcal{A}$, $i = 1, \dots, p$ such that $\alpha_i \in A_i$. Choosing r_i sufficiently close to $x(\alpha_i)$, we obtain $x \in C_\sigma$, where $\sigma = \{(A_1, \dots, A_p), (r_1, \dots, r_p)\}$.

Let us estimate the diameter of C_σ . If $x, y \in C_\sigma$, choose $\alpha_i \in A_i$, $\beta_i \in A_i$, ($i = 1, \dots, p$) such that

$$\sum_{i=1}^p (x(\alpha_i) - r_i)^2 < \varepsilon^2 \quad \text{and} \quad \sum_{i=1}^p (y(\beta_i) - r_i)^2 < \varepsilon^2.$$

Put $x_1(\alpha_i) = x(\alpha_i)$, $x_2(\alpha_i) = r_i$ for $i = 1, \dots, p$,

$$x_1(\alpha) = x_2(\alpha) = 0 \text{ for all other } \alpha,$$

$$y_1(\beta_i) = y(\beta_i), \quad y_2(\beta_i) = r_i \text{ for } i = 1, \dots, p,$$

$$y_1(\beta) = y_2(\beta) = 0 \text{ for all other } \beta.$$

Then $1 = \|x - x_1\|^2 + \|x_1\|^2 \geq \|x - x_1\|^2 + (\|x_2\| -$

$$- \|x_1 - x_2\|)^2 \geq \|x - x_1\|^2 + (1 - 2\varepsilon)^2$$

thus $\|x - x_1\|^2 \leq 4\varepsilon - 4\varepsilon^2 \leq 4\varepsilon;$

similarly we prove that $\|y - y_1\| \leq 2\sqrt{\varepsilon}$, therefore
 $\|x - y\| \leq \|x - x_1\| + \|x_1 - x_2\| + \|x_2 - y_2\| + \|y_2 - y_1\| + \|y_1 - y\| \leq \sqrt{2} + 4\sqrt{\varepsilon} + 2\varepsilon$.

(iv) \Rightarrow (iii): We can suppose that $m = 2^n$ and n is a set of ordinals such that $\text{card } T_\alpha < n$ for any $\alpha \in n$. For $\alpha \in n$ and $B \subset T_\alpha$ put $A_{\alpha, B} = \{C \subset n; C \cap T_\alpha = B\}$. The family $\{A_{\alpha, B}; \alpha \in n, B \subset T_\alpha\}$ separates points in 2^n and, since $2^{\text{card } T_\alpha} \leq n$, its cardinality is $\leq n$.

Remark 1: Not using the continuum hypothesis we can prove (in the same way as in (iv) \Rightarrow (iii)) that (iii) holds for such cardinals n, m that

- (a) $m \leq 2^n$
- (b) If $n' < n$ then $2^{n'} \leq n$.

Remark 2: If $\aleph_\alpha \leq n < m$ are cardinal numbers satisfying the condition (iii) of the theorem and if $n^{\aleph_\alpha} = n$ then the unit sphere in $\ell_2(m)$ can be written as a union of n sets with diameter $\leq \sqrt{2}$. (One can take the covers \mathcal{C}_ρ with diameter $< \sqrt{2} + \frac{1}{\rho}$ and put $\mathcal{C} = \bigcup_{\rho=1}^{\infty} A_{n, \rho}$; $A_{n, \rho} \in \mathcal{C}_\rho$.) Therefore the graphs obtained by joining two points of the $\aleph_{\alpha+2}$ -dimensional Hilbert space if their distance is $> \sqrt{2}$ has the chromatic number $\aleph_{\alpha+1}$.

R e f e r e n c e

- [1] P. ERDÖS and A. HAJNAL: On chromatic number of graphs and set-systems, Acta Math. Acad. Sci. Hung. 17 (1966), 61-99.

Mémetvölgyi ut 72/c
1124 Budapest
Hungary

Matematicko-fyzikální fakulta
Karlova universita
Sokolovská 83, 18600 Praha 8
Československo

(Oblatum 9.9. 1976)

