Werk

Label: Errata

Jahr: 1979

PURL: https://resolver.sub.uni-goettingen.de/purl?316342866_0020|log21

Kontakt/Contact

Digizeitschriften e.V.
SUB Göttingen
Platz der Göttinger Sieben 1
37073 Göttingen

info@digizeitschriften.de

COMMENTATIONES MATHEMATICAE UNIVERSITATIS CAROLINAE
20, 1 (1979)

CORRIGENDUM ET ADDENDUM AD "MINIMAL CELL COVERINGS OF SPHERE BUNDLES OVER SPHERES"

D. A. MORAN

It has been pointed out by Prof. Nelson Max [3] that the key step in the purported proof of the main theorem in [5] is in error. I have been unable to recover the full strength of that result, but I wish to delineate the circumstances wherein it has been proved. The terminology and notation of [5] will be used.

1. An easy argument employing the exact homotopy sequence of a bundle shows that M is k-connected, where k = = min (p,q) - 1. According to theorems of Luft [2] and of Osborne and Stern [6], it can be inferred from this that M can be covered by three cells if $\frac{1}{2}(p+1) \leq q \leq 2p-1$.

2. By Bott's famous computations, if $p \equiv 3$, 5 or 6 (mod 8) and $q + 2 > p$, then $\pi_{p-1}(SO_{q+1}) = 0$. For such p and q, all q-sphere bundles over a p-spheres are products, and can be covered by three cells.

3. If the fibration admits a global cross-section, M can be covered by three cells [4].

To my knowledge, the remaining cases are still open.

References

[1] R. BOTT: The stable homotopy of the classical groups, Ann. of Math. 70(1959), 313-337.

[2] E. LUFT: Covering of manifolds with open cells, Illinois J. Math. 13(1969), 321-326.

[3] N.L. MAX: Math. Reviews, vol. 51 # 1834.

[4] D. MORAN: Minimal cell coverings of some sphere bundles, Comment. Math. Univ. Carolinae 14(1973), 647-650.

[5] D. MORAN: Minimal cell coverings of sphere bundles over spheres, Comment. Math. Univ. Carolinae 16(1975), 147-150.

[6] R. OSBORNE and J. STERN: Covering manifolds with cells, Pacific J. Math. 30(1969), 201-207.

Michigan State University

East Lansing, MI 48824

U.S.A.

(Oblatum 30.11. 1978)