

## Werk

**Label:** Article

**Jahr:** 1979

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?316342866\\_0020|log44](https://resolver.sub.uni-goettingen.de/purl?316342866_0020|log44)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

ON DOMINATION PROBLEM IN BANACH ALGEBRAS  
Vladimir MULLER

**Abstract:** We give an example of a commutative Banach algebra  $A$  with elements  $u, v, w \in A$  such that  $|ux| \leq |vx| + |wx|$  for every  $x \in A$  and there exists no commutative Banach algebra  $B$  containing  $A$  as a subalgebra and elements  $b, c \in B$  such that  $u = bv + cw$ . This gives the negative answer to the problem of Zelazko [4].

**Key words:** Banach algebras, ideals.

AMS: 46H05

---

**Introduction.** Let  $A$  be a unital commutative complex Banach algebra, let  $u, v_1, \dots, v_n$  ( $1 \leq n < \infty$ ) be elements of  $A$ .

As in [4] we say that  $u$  is dominated by elements  $v_1, \dots, v_n$  if there exists a constant  $k \geq 0$  such that  $|ux| \leq k \cdot \sum_{i=1}^n |v_i x|$  for every  $x \in A$ .

Let  $A, B$  be unital commutative complex Banach algebras. We say that  $B$  is an isometric extension of  $A$  if there exists a unit preserving isometric isomorphism from  $A$  into  $B$ . In this case we consider  $A$  as a subalgebra of  $B$  and write  $A \subset B$ .

Let  $A \subset B$ ,  $u, v_1, \dots, v_n \in A$ . Let  $u = \sum_{i=1}^n b_i v_i$  for some  $b_i \in B$ . Then  $|ux| \leq \sum_{i=1}^n |b_i| |v_i x| \leq k \cdot \sum_{i=1}^n |v_i x|$  for each  $x \in A$ , where  $k = \max(|b_1|, \dots, |b_n|)$ . So  $u$  is dominated by the elements  $v_1, \dots, v_n$ .

In [4] (Problem 9), the question was raised whether the converse statement is true. More precisely:

Let  $u, v_1, \dots, v_n \in A$ , let  $u$  be dominated by  $v_1, \dots, v_n$ . Does it follow that in some isometric extension  $B \supset A$  there are elements  $b_1, \dots, b_n$  such that  $u = \sum_{i=1}^n b_i v_i$ ? The answer is positive for  $n = 1$  ([1]) and also for arbitrary  $n$  in special Banach algebras ([5]). In the present paper we give an example that this is not true for  $n = 2$  (and of course for  $n \geq 2$ ) in general Banach algebras.

Lemma. There exists a unital commutative complex Banach algebra  $A$  satisfying the following conditions:

- 1) There are (distinct) elements  $u, v, w, a_{ij}$  ( $i, j = 0, 1, 2, \dots$ ) in  $A$  generating  $A$ .
- 2)  $u^2 = v^2 = w^2 = uv = uw = vw = 0, a_{ij}a_{km} = 0$  for every  $i, j, k, m \geq 0$
- 3)  $a_{ij}u = a_{i-1, j}v + a_{i, j-1}w$  ( $i, j \geq 1$ )  
 $a_{i, 0}u = a_{i-1, 0}v$  ( $i \geq 1$ )  
 $a_{0, j}u = a_{0, j-1}w$  ( $j \geq 1$ )
- 4)  $|a_{ij}| = 2^{-(i+j)^2}$  ( $i, j \geq 0$ ),  $|a_{0, 0}u| = 1$
- 5)  $u$  is dominated by  $v, w$ .

Construction: Let  $S$  be the free commutative semigroup with unit 1 and zero element 0 ( $0s = 0$  for each  $s \in S$ ) and with generators  $u', v', w', a'_{ij}$  ( $i, j = 0, 1, 2, \dots$ ) satisfying  $u'^2 = v'^2 = w'^2 = u'v' = u'w' = v'w' = 0, a'_{ij}a'_{km} = 0$  ( $i, j, k, m = 0, 1, 2, \dots$ ). Put  $|u'| = |v'| = |w'| = 1, |a'_{ij}| = |a'_{ij}u'| = |a'_{ij}v'| = |a'_{ij}w'| = 2^{-(i+j)^2}$  for every  $i, j \geq 0$ .

Let  $B$  be the  $\ell^1$  algebra over semigroup  $S$  with this norm,

i.e. B is formed by formal linear combinations

$$(1) \quad x = \lambda_0 + \lambda_1 u' + \lambda_2 v' + \lambda_3 w' + \sum_{i,j=0}^{\infty} \lambda_{ij} a'_{ij} + \\ + \sum_{i,j=0}^{\infty} (\mu_{ij}^{(1)} a'_{ij} u' + \mu_{ij}^{(2)} a'_{ij} v' + \mu_{ij}^{(3)} a'_{ij} w')$$

where  $\lambda_0, \dots, \lambda_3, \lambda_{ij}, \mu_{ij}^{(k)}$  ( $k = 1, 2, 3, i, j = 0, 1, \dots$ ) are complex numbers and

$$|x| = \sum_{i=0}^3 |\lambda_i| + \sum_{i,j=0}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2} + \\ + \sum_{k=1}^3 \sum_{i,j=0}^{\infty} |\mu_{ij}^{(k)}| 2^{-(i+j)^2} < \infty.$$

Clearly B with this norm is a unital commutative Banach algebra. Let  $I \subset B$  be the closed ideal generated by elements

$$a'_{ij} u' - a'_{i-1,j} v' - a'_{i,j-1} w' \quad (i, j \geq 1),$$

$$a'_{i,0} u' - a'_{i-1,0} v' \quad (i \geq 1) \text{ and } a'_{0,j} u' - a'_{0,j-1} w' \quad (j \geq 1).$$

Denote  $A = B/I, u = u' + I, v = v' + I, w = w' + I, a_{ij} = a'_{ij} + I$  ( $i, j = 0, 1, \dots$ ). We prove that A satisfies all the conditions required.

Conditions 1), 2) and 3) are trivial.

Let us notice that if  $x \in B, x$  has the form (1), then

$$(2) \quad |x + I|_A = \sum_{i=0}^3 |\lambda_i| + \sum_{i,j=0}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2} + |\mu_{00}^{(1)}| + \\ + \sum_{i+j \geq 1} \inf_{\lambda \in \mathbb{C}} |\mu_{ij}^{(1)} a'_{ij} u' + \mu_{i-1,j}^{(2)} a'_{i-1,j} v' + \mu_{i,j-1}^{(3)} a'_{i,j-1} w' + \\ + \nu a'_{ij} u' - \nu a'_{i-1,j} v' - \nu a'_{i,j-1} w'|_B = \sum_{i=0}^3 |\lambda_i| + \\ + \sum_{i,j=0}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2} + \sum_{i,j=0}^{\infty} |\mu_{ij}^{(1)} a'_{ij} u' + \mu_{i-1,j}^{(2)} a'_{i-1,j} v' + \\ + \mu_{i,j-1}^{(3)} a'_{i,j-1} w'|_A.$$

(Here we put  $a_{km} = a'_{km} = 0$  for  $\min(k, m) < 0$ .)

From formula (2), the condition 4) immediately follows.

Further, it holds

$$\begin{aligned}
 (3) \quad |a_{ij}u| &= 2^{-(i+j)^2} \quad (i, j \geq 0) \\
 |a_{ij}v| &= 2^{-(i+j)^2} \quad (i \geq 0, j \geq 1) \\
 |a_{ij}w| &= 2^{-(i+j)^2} \quad (i \geq 1, j \geq 0) \\
 |a_{i,0}v| &= |a_{i+1,0}u| = 2^{-(i+1)^2} \quad (i \geq 0) \\
 |a_{0,j}w| &= |a_{0,j+1}u| = 2^{-(j+1)^2} \quad (j \geq 0).
 \end{aligned}$$

It remains to prove the condition 5). Let  $x \in B$  have the form (1),  $y = x + I \in A$ . Then by (2), (3)

$$\begin{aligned}
 |yu|_A &= |xu + I|_A = |\lambda_0 u' + \sum_{i,j=0}^{\infty} \lambda_{ij} a'_{ij} u' + I|_A = |\lambda_0| + \\
 &\quad + \sum_{i,j=0}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2}, \\
 |yv|_A &= |xv' + I|_A = |\lambda_0 v' + \sum_{i,j=0}^{\infty} \lambda_{ij} a'_{ij} v' + I|_A = |\lambda_0| + \\
 &\quad + \sum_{i=0}^{\infty} \sum_{j=1}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2} + \sum_{i=0}^{\infty} |\lambda_{i,0}| 2^{-(i+1)^2}, \\
 |yw|_A &= |xw' + I|_A = |\lambda_0 w' + \sum_{i,j=0}^{\infty} \lambda_{ij} a'_{ij} w' + I|_A = |\lambda_0| + \\
 &\quad + \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} |\lambda_{ij}| 2^{-(i+j)^2} + \sum_{j=0}^{\infty} |\lambda_{0,j}| 2^{-(j+1)^2}.
 \end{aligned}$$

From this immediately follows  $|yu| \leq |yv| + |yw|$  for every  $y \in A$ , hence  $u$  is dominated by  $v, w$ . Note that  $u$  is not dominated by  $v$ . It is  $|a_{i,0}u|/|a_{i,0}v| = 2^{-i^2}/2^{-(i+1)^2} = 2^{2i+1}$  which forms an unbounded sequence. Similarly, neither  $u$  is dominated by  $w$ .

**Theorem.** Let  $A$  be the Banach algebra from the previous Lemma. Let  $C$  be any isometric extension of  $A$ . Then there

exist no  $b, c \in C$  such that  $u = bv + cw$ .

Proof: I. Let  $k > 0$  be fixed. Let  $B_k$  be the  $\ell^1$  algebra over the free commutative semigroup with generators  $b_k, c_k$  with coefficients in  $A$ , i.e.  $B_k$  consists of elements of the form  $x = \sum_{i,j=0}^{\infty} x_{ij} b_k^i c_k^j$ , where  $x_{ij} \in A$  ( $i, j \geq 0$ ) and  $\|x\|_{B_k} = \sum_{i,j=0}^{\infty} |x_{ij}|_A \cdot k^{i+j}$ .

Algebraic operations in  $B_k$  are defined as follows:

For  $y = \sum_{i,j=0}^{\infty} y_{ij} b_k^i c_k^j$  it is  $x + y = \sum_{i,j=0}^{\infty} (x_{ij} + y_{ij}) b_k^i c_k^j$ ,

$xy = yx = \sum_{m,n=0}^{\infty} b_k^m c_k^n (\sum_{\substack{i+j=m \\ i'+j'=n}} x_{ij} y_{i'j'})$ .

Clearly  $B_k$  is a Banach algebra,  $B_k \supset A$ . Denote  $z = u - b_k v - c_k w$ . Let  $J = \overline{zB_k}$  be the closed ideal generated by  $z$ . Denote  $d = \sum_{i,j=0}^{\infty} a_{ij} b_k^i c_k^j$  where  $a_{ij}$  are elements from the previous Lemma. It holds

$$\|d\|_{B_k} = \sum_{i,j=0}^{\infty} |a_{ij}|_A \cdot k^{i+j} = \sum_{i,j=0}^{\infty} 2^{-(i+j)^2} \cdot k^{i+j} = \sum_{m=0}^{\infty} 2^{-m^2}.$$

$\cdot k^{m(m+1)} < \infty$ . So  $d \in B_k$ . We have

$$\begin{aligned} dz &= (\sum_{i,j=0}^{\infty} a_{ij} b_k^i c_k^j)(u - b_k v - c_k w) = a_{0,0} u + \sum_{j=1}^{\infty} b_k^j (a_{1,0} u - \\ &- a_{i-1,0} v) + \sum_{j=1}^{\infty} c_k^j (a_{0,j} u - a_{0,j-1} w) + \sum_{i,j=1}^{\infty} b_k^i c_k^j (a_{ij} u - \\ &- a_{i-1,j} v - a_{i,j-1} w) = a_{0,0} u. \text{ Hence } a_{0,0} u \in J. \end{aligned}$$

II. Suppose now on the contrary that there exists a Banach algebra  $C$  containing  $A$  as a subalgebra and  $b, c \in C$  such that  $u = bv + cw$ . Choose  $k \geq \max(|b|, |c|)$ . Define a homomorphism  $f: B_k \rightarrow C$  by  $f(\sum_{i,j=0}^{\infty} x_{ij} b_k^i c_k^j) = \sum_{i,j=0}^{\infty} x_{ij} b^i c^j$ . It is

$$\begin{aligned} \|\sum_{i,j=0}^{\infty} x_{ij} b^i c^j\|_C &\leq \sum_{i,j=0}^{\infty} |x_{ij}|_A |b|^i |c|^j \leq \sum_{i,j=0}^{\infty} |x_{ij}|_A \cdot k^{i+j} = \\ &= \|\sum_{i,j=0}^{\infty} x_{ij} b_k^i c_k^j\|_{B_k}. \end{aligned}$$

So the definition of  $f$  is correct and  $\|f\| \leq 1$ . Clearly

$f(b_k) = b$ ,  $f(c_k) = c$  and  $f|_A$  is the identical mapping (we identify elements of  $A$  with the corresponding elements of  $B_k$  and  $C$ , respectively). It holds  $f(z) = f(u - b_k v - c_k w) = u - b v - c w = 0$ , so  $f(J) = 0$ . Hence  $f(a_{0,0} u) = 0$ . On the other hand,  $a_{0,0} u \in A$  and  $f|_A$  is the identical mapping. Necessarily  $a_{0,0} u = 0$  which contradicts the condition 4) of Lemma.

Remark 1: A Banach algebra  $B$  is called an extension of a Banach algebra  $A$  if there exists a unit preserving topological isomorphism of  $A$  into  $B$ . It is easy to see that the words "isometric extension" in the Theorem can be replaced by "extension". The proof in this case is the same. Note also that every extension  $C$  of  $A$  becomes an isometric extension after a suitable renorming of  $C$  (see [2]).

Remark 2: The following question still remains open: Let  $1$  (unit element of  $A$ ) be dominated by  $v_1, \dots, v_n \in A$ . Does it follow that  $1 = \sum_{i=1}^n b_i v_i$  for some extension  $B$  of  $A$  and some  $b_i \in B$ ?

This question is equivalent to Problem 5 of [4]: Does every non-removable ideal in  $A$  consist of joint topological divisors of zero?

For related topics see also [3].

#### R e f e r e n c e s

- [1] R. ARENS: Extensions of Banach algebras, Pacific J. Math. 10(1960), 1-16.
- [2] J.A. LINDBERG: Extensions of algebra norms and applications, Studia Math. 40(1971), 35-39.
- [3] W. ZELAZKO: Banach algebras, Elsevier-Publishing Company (Polish Scientific Publishers, Amsterdam-London-

New York) Warszawa, 1973.

- [4] W. ZELAZKO: On some classes of ideals in commutative Banach algebras, Rend. Sem. Mat. Fis. Milano (in print).
- [5] W. ZELAZKO: A domination theorem for function algebras (to appear).

Matematický ústav ČSAV  
Žitná 25, 11567 Praha 1  
Československo

(Oblatum 23.2. 1979)



