

## Werk

**Label:** Article

**Jahr:** 1982

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?316342866\\_0023|log56](https://resolver.sub.uni-goettingen.de/purl?316342866_0023|log56)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

RESTRICTED MEAN VALUE PROPERTY IN AXIOMATIC  
POTENTIAL THEORY  
Jiří VESELY

**Abstract:** A generalization of the restricted mean value property and a corresponding converse of the Gauss mean value theorem are studied in the frame of the axiomatic potential theory.

**Key words:** Mean value property, axiomatic potential theory, Dirichlet problem,  $\mathcal{O}$ -harmonicity.

**Classification:** Primary 31D05, 31C05  
Secondary 31A25

---

**Introduction.** Let  $G \subset \mathbb{R}^n$  be an open set having non-empty boundary  $\partial G$  and let  $\mathcal{O}$  be an arbitrary function on  $G$  such that  $0 < \mathcal{O}(x) < \text{dist}(x, \partial G)$ , the distance of  $x$  from the boundary. A Borel measurable function  $g$  on  $G$  is said to have the restricted mean value property with respect to balls, if for every  $x \in G$  the mean value of  $g$  over the ball with centre  $x$  and radius  $\mathcal{O}(x)$  (i.e. the integral of  $g$  over the ball divided by its volume) is equal to  $g(x)$ . A function  $g$  with the described property is sometimes called  $\mathcal{O}$ -harmonic function on  $G$ . Not so frequent is the study of  $\mathcal{O}$ -harmonicity with respect to spheres where the mean value may be interpreted as the integral with respect to harmonic measure, nevertheless this serves to us as a starting point for our generalization of the problem.

A quite natural generalization in the context of a harmonic space  $X$  may be formulated as follows: For every  $x$  from an open set  $G \subset X$  fix a measure  $\nu_x$  such that for every function  $h \in \mathcal{H}(G)$ , the set of harmonic functions on  $G$ , we have

$$\nu_x(h) = \int h d\nu_x = h(x).$$

Now the problem consists of finding conditions involving  $G$ ,  $g$  and the choice of measures  $\nu_x$  which imply  $g \in \mathcal{H}(G)$  provided  $\nu_x(g) = g(x)$  for every  $x \in G$ . A lot of results is known for the classical case and various conditions imposed on  $G$ ,  $g$  and  $\sigma$  which give the validity of

$$g \text{ is } \sigma\text{-harmonic} \implies g \in \mathcal{H}(G)$$

were studied (see, e.g. [5],[12],[4]; a survey article [9] contains 81 references concerning the problem). As far as we know in the frame of an axiomatics such a problem has not been studied. Our approach to it is closely connected with the Dirichlet problem and avoids probabilistic methods which occurred to be very efficient in the classical case. Its origin goes back to Lebesgue [7] (this influenced our terminology). Other comments may be found at the end of the article.

1. Notation. Let  $X$  be a  $\mathcal{P}$ -harmonic space with countable base in the sense of [3]. In what follows,  $G \subset X$  is a fixed open set for which  $\partial G \neq \emptyset$ . For a set  $M \subset X$ , the symbol  $\mathcal{C}(M)$  stands for the set of continuous functions on  $M$  while  $\mathcal{B}(M)$  denotes the set of Borel measurable functions on  $M$ . The set of functions from  $\mathcal{C}(X)$  having compact support is denoted by  $\mathcal{C}_c(X)$  and  $\mathcal{B}_b(M)$  is the subset of  $\mathcal{B}(M)$  consisting of bound-

ed functions.

For an open relatively compact  $G \subset X$  let us denote by  $S(\bar{G}) = S$  the cone of lower semicontinuous functions (l.s.c.) on  $\bar{G}$  which are extensions of superharmonic functions on  $G$ . For every  $x \in \bar{G}$  choose a representing measure  $\alpha_x$  with respect to  $S$ , i. e. a measure  $\alpha_x$  such that

$$(1) \quad s \in S \Rightarrow \alpha_x(s) \leq s(x).$$

Redefining for a function  $s \in S$  the value  $s(x)$  for an  $x \in \partial G$  in such a way that  $s(x) < s(y)$  for all  $y \in \bar{G} \setminus \{x\}$  we can easily show  $\alpha_x = \epsilon_x$  (the Dirac measure); hence  $\alpha_x = \epsilon_x$  for all  $x \in \partial G$ .

Now we can put for  $f \in \mathcal{B}_b(G)$

$$(2) \quad Af(x) := \alpha_x(f)$$

where we identify  $A$  with the collection  $\{\alpha_x\}_{x \in \bar{G}}$ . Since mostly the choice of  $\alpha_x$  together with (2) does not determine a "good operator"  $A$ , we must be a bit careful:  $Af = f$  means only that  $f$  is  $\alpha_x$ -integrable for all  $x$  and

$$(3) \quad \alpha_x(f) = f(x).$$

If (2) determines an operator acting on the space  $\mathcal{C}(\bar{G})$  or  $\mathcal{B}_b(\bar{G})$ , we shall call  $A$  "continuous" or "Borel" operator, respectively. Our first aim will be to find sufficient conditions for the validity of

$$(4) \quad Af = f \Rightarrow f|G \in \mathcal{H}(G);$$

here  $f|G$  means the restriction of  $f$  on  $G$ . Then we shall use it in connection with the restricted mean value property.

2. Lebesgue's operators. Let  $G$  be a relatively compact open set. For  $A = \{\alpha_x\}_{x \in \bar{G}}$  suppose (1) and

$$(5) \quad \alpha_x \neq \epsilon_x \text{ for all } x \in G.$$

In what follows we shall need a variant of the minimum principle:

If  $S_A$  is the set of l.s.c. functions  $s$  on the compact set  $\bar{G}$  for which  $As \leq s$ , then for every  $s \in S_A$  we have

$$(6) \quad s|_{\partial G} \geq 0 \Rightarrow s \geq 0.$$

Suppose, on the contrary, that there is a function  $s \in S_A$  such that  $s(x) < 0$  for an  $x \in G$  and  $s|_{\partial G} \geq 0$ . If  $p > 0$  is a continuous potential on  $X$ , put  $f = s$ ,  $g = p|_{\bar{G}}$  in Lemma 2 ([3], p. 26) to obtain a contradiction with (5). By a Lebesgue's operator  $A = \{\alpha_x\}_{x \in \bar{G}}$  (shortly: L-operator) we understand a continuous operator defined by (2) provided (1) and (5) hold. Denote by  $\alpha_x^n$  the measure determined for all  $x$  and  $n$  with the help of the equality

$$(7) \quad \alpha_x^n(f) := A^n f(x), \quad f \in \mathcal{C}(\bar{G})$$

(here  $A^0$  is the identity operator and  $A^n = A * A^{n-1}$ ,  $n = 1, 2, \dots$ , where  $*$  stands for the composition of operators) and by  $\mu_x$  the harmonic measure corresponding to  $G$  and  $x$ .

3. Proposition. Let  $G \subset X$  be an open relatively compact set and  $A = \{\alpha_x\}_{x \in \bar{G}}$  be an L-operator. If  $\alpha_x^n$  are defined by (7), then for every  $f \in \mathcal{C}(\bar{G})$

$$(8) \quad \alpha_x^n(f) \rightarrow \mu_x(f), \quad x \in G$$

as  $n \rightarrow \infty$  (i.e.  $\alpha_{\mathbf{x}}^n \rightarrow \mu_{\mathbf{x}}$  weakly) and the convergence in (8) is uniform with respect to  $\mathbf{x}$  on compact subsets of  $G$ .

**Proof.** Since the differences of continuous potentials are uniformly dense in  $\mathcal{C}_0(X)$  (cf. [3], Th. 2.3.1), we restrict ourselves to the case of  $f \in \mathcal{C}(\bar{G})$  and  $f|_G$  superharmonic. Then we have

$$f_1 := Af \leq f, \quad f_2 := Af_1 \leq f_1, \dots$$

and  $\{f_n\}$  is lower bounded. Put  $u = \lim f_n$ . It is easily seen that  $v(\mathbf{x}) := \mu_{\mathbf{x}}(f) \leq u(\mathbf{x})$ ,  $\mathbf{x} \in G$  and  $Au = u$ .

If  $s$  is an upper function for  $f|_{\partial G}$ , we extend the difference  $(s-u)$  by  $\liminf$  from  $G$  onto  $\bar{G}$ . Then  $s-u \in S_A$  and by (6) we have  $u \leq s$ . Consequently  $u = v$ . The rest follows from the Dini's theorem (cf. [13], Th.6).

**4. Corollary.** Let  $G \subset X$  be an open relatively compact set. Suppose that for a Borel operator  $A = \{\alpha_{\mathbf{x}}\}_{\mathbf{x} \in \bar{G}}$  there is an L-operator  $B = \{\beta_{\mathbf{x}}\}_{\mathbf{x} \in \bar{G}}$  for which

$$(9) \quad s \in S \Rightarrow \alpha_{\mathbf{x}}(s) \leq \beta_{\mathbf{x}}(s), \quad \mathbf{x} \in G.$$

Then (8) is valid for every  $f \in \mathcal{C}(\bar{G})$ .

Indeed, it is easily seen that for every  $s \in S$  we have  $\mu_{\mathbf{x}}(s) \leq \alpha_{\mathbf{x}}^n(s) \leq \beta_{\mathbf{x}}^n(s) \rightarrow \mu_{\mathbf{x}}(s)$  and hence the statement is an easy consequence of (9). In connection with the Dirichlet problem both previous assertions offer the solution as a limit of a sequence of functions. More precisely, if  $f \in \mathcal{C}(\bar{G})$  is an extension of the boundary condition in the Dirichlet problem and  $A$  is as above, then  $A^n f$  is convergent on  $G$  to the solution of the problem (see [13]).

Now we shall turn to the restricted mean value property. We shall suppose

$$(10) \quad \text{spt } \alpha_x \subset G \text{ for all } x \in G;$$

this assumption is quite natural with regard to the classical case we discussed in the introduction. The same may be said on the definition:

If  $G \subset X$  is open and  $A = \{\alpha_x\}_{x \in G}$  is a collection of measures having compact support and satisfying (1), (5) and (10), then a locally bounded  $f \in \mathcal{B}(G)$  is said to be  $A$ -harmonic on  $G$  provided (3) holds for every  $x \in G$ .

5. Proposition. Suppose that  $G \subset X$  is an open relatively compact set and  $A = \{\alpha_x\}_{x \in G}$  is a Borel operator (i. e.  $Ag \in \mathcal{B}_b(G)$  whenever  $g \in \mathcal{B}_b(G)$ ) for which there is an  $L$ -operator  $B = \{\beta_x\}_{x \in \bar{G}}$  such that (9) is valid. If  $f \in \mathcal{B}_b(\bar{G})$  and  $f$  is  $\mu_x$ -almost everywhere continuous on  $\bar{G}$  for every  $x \in G$ , then from  $A$ -harmonicity of  $f$  follows  $f|_G \in \mathcal{H}_b(G)$ .

This is a consequence of Cor. 4, namely the weak convergence  $\alpha_x^n \rightarrow \mu_x$ , which we established there and from which it follows for  $f$

$$f(x) = \alpha_x^n(f) = \mu_x(f), \quad x \in G.$$

(See [1], Th. 4.5.1.)

The condition (9) shows that for every  $x \in G$  the corresponding  $\alpha_x$  is balayed  $\beta_x$  with respect to the cone  $S$ . Roughly speaking,  $A$ -harmonicity implies harmonicity if  $A$  is a Borel operator (on  $\mathcal{B}_b(G)$ ) which is a "pointwise balayed  $L$ -operator".

It seems a bit inconvenient to work with an  $f$  defined on

$\bar{G}$  since the result is naturally connected with its values on  $G$ . This is removed by the next theorem.

6. Theorem. Let  $G \subset X$  be an open relatively compact set. Suppose that  $f \in \mathcal{B}_b(G)$  is an  $A$ -harmonic function on  $G$  with respect to the Borel operator  $A = \{\alpha_x\}_{x \in G}$ . Moreover, suppose that (i) there is an  $L$ -operator  $B = \{\beta_x\}_{x \in \bar{G}}$  and (9) holds, (ii) there is a Borel set  $N_f \subset \partial G$  such that

$$(11) \quad \lim_{x \rightarrow y} f(x)$$

is finite for every  $y \in \partial G \setminus N_f$  and (iii) the set  $N_f$  is negligible in the sense that

$$(12) \quad \mu_x(N_f) = 0 \text{ for every } x \in G.$$

Then  $f$  is a bounded harmonic function on  $G$ .

Proof. Consider the weak convergence  $\alpha_x^n \rightarrow \mu_x$  in (8) for a fixed  $x \in G$ . Recall that by (10)  $\text{spt } \alpha_x^n \subset G$  while for the harmonic measure we have  $\text{spt } \mu_x \subset \partial G$ . Put  $P = \bar{G} \setminus N_f$ ,  $\alpha_x^n|_P = \alpha_n$ ,  $\mu_x|_P = \gamma$ . It is easily seen that the function  $f$  extended from  $G$  on  $P$  by limits (11) is a Borel measurable function which is  $\gamma$ -almost everywhere continuous on  $P$ . It remains to prove  $\alpha_n \rightarrow \gamma$  weakly and again to apply Th. 4.5.1 in [1], where also equivalent conditions for the weak convergence of measures may be found. For any  $V \subset P$  open there is an open  $V' \subset \bar{G}$  such that  $V' \cap P = V$ . Since we know (8), we can easily prove  $\liminf_{n \rightarrow \infty} \alpha_n(V) \geq \gamma(V)$  and the required weak convergence will be established. Indeed,  $\liminf \alpha_n(V) = \liminf \alpha_x^n(V) = \liminf \alpha_x^n(V') \geq \mu_x(V') = \mu_x(V) = \gamma(V)$ , and hence  $\alpha_n \rightarrow \gamma$  is proved.



Let us remark that it is not obvious whether (12) is equivalent with  $\nu(N_f) = 0$  for a measure  $\nu$ . But it is the case if we choose a countable dense subset  $\{x_n; n \in \mathbb{N}\} \subset G$  and put  $\nu = \sum_{n=1}^{\infty} 2^{-n} \mu_{x_n}$ . Since there is a continuous potential  $p > 0$  on  $X$ , it is easily seen that the measures  $\mu_{x_n}$  are uniformly bounded and hence  $\nu$  is a finite measure. Now, if (12) holds,  $\nu(N_f) = 0$  is obvious. On the other hand, if  $\nu(M) = 0$  for an  $M \subset \partial G$ , then  $\mu_{x_n}(M) = 0$  for every  $n \in \mathbb{N}$ . Since  $\mu_x(M)$  is - as a function of  $x$  on  $G$  - harmonic and therefore continuous on  $G$ , it is equal 0 everywhere and (12) is valid.

To find a connection of the theorem to the results known in the classical case, we shall at first examine collections  $A$  of "typical"  $\alpha_x$ . If  $V_x$  is a neighbourhood of  $x \in G$  such that  $\overline{V_x} \subset G$  is compact, then a good candidate for such an  $\alpha_x$  is  $\varepsilon_x^{CV}$  (the harmonic measure corresponding to the set  $V_x$  at  $x$ ). Since the support of the measure is compact, every  $f \in \mathfrak{B}_p(G)$  will be  $\alpha_x$ -integrable. In what follows,  $U, V$  (sometimes with primes or indexes) will stand for open relatively compact subsets of  $G$ . Let us fix a metric  $\varphi$  on  $X$  compatible with its topology and introduce

$$\mathcal{G}(U, V) = \tilde{\varphi}(\overline{U}, \overline{V}) + \tilde{\varphi}(\partial U, \partial V),$$

where  $\tilde{\varphi}$  is the Hausdorff metric on the system of compact subsets of  $X$ . It is easily seen that  $\mathcal{G}$  is a metric on the set of open relatively compact subsets of  $G$ . Moreover, if  $\varepsilon > 0$  is given and for a  $U$  we put

$$V' = \{x \in U; \text{dist}(x, \partial U) > \varepsilon\}, \quad V'' = \{x \in G; \text{dist}(x, U) < \varepsilon\},$$

then  $V' \subset U \subset V''$  and for every  $V$ ,  $\mathcal{G}(U, V) < \varepsilon$  we have

$V' \subset V \subset V''$ . For a technical reason let us put

$$\alpha(U, x, f) := \int f d\epsilon_x^{CU}$$

for  $x \in U$ ,  $\bar{U} \subset G$  and  $f \in \mathcal{B}_b(G)$ .

**7. Proposition.** Let  $G \subset X$  be an open set. Suppose that the mapping  $\omega: x \rightarrow V_x$ ,  $x \in G$  has the following properties:

(i)  $V_x$  is a neighbourhood of  $x$  such that  $\bar{V}_x \subset G$  is compact and  $\epsilon_x^{CV} = \epsilon_x^{C\bar{V}}$ ,

(ii)  $\omega$  is continuous (w.r.t.  $\varphi - \epsilon$ ).

Then, for every  $f \in \mathcal{C}(G)$ , the function

$$Af(x) := \alpha(V_x, x, f), \quad x \in G$$

is continuous (here  $\alpha(V_x, x, \cdot)$  denotes a measure previously denoted by  $\alpha_x$ ).

**Remark.** The equality of measures in (i) implies certain continuity: put  $V = V_x$  and choose  $U_n, U'_n$  in such a way that  $U_n \nearrow V, U'_n \searrow V$ . Then it follows from Pr. 7.2.4 in [3] that  $\epsilon_y^{CV_n} \rightarrow \epsilon_y^{CV}, \epsilon_y^{CV'_n} \rightarrow \epsilon_y^{C\bar{V}}$  for every  $y \in V$  and hence our assumption implies that both limits coincide for  $y = x$ . (The condition is fulfilled if the set of points of  $\partial V$  in which  $\bar{C}\bar{V}$  is thin is "small"; cf. [10], Th. 1.)

**Proof.** Let us prove the continuity of  $Af$  at arbitrarily chosen  $x \in G$ . For such a fixed  $x$  choose  $U, \bar{V}_x \subset U \subset \bar{U} \subset G$ . Now if  $\sigma(V_x, V_y)$  is sufficiently small, then  $\bar{V}_y \subset U$  and hence the values  $Af(y)$  depend only on values of  $f$  on the set  $\bar{U}$ . From this it follows with the help of approximation theorem Th. 2.3.1 in [3] that it is enough to prove the continuity of  $A_p$  at  $x$  for any potential  $p$  on  $X$ .

The assumption (i) implies (see the remark above) the existence of neighbourhoods  $U, U'$  of  $x$  such that  $\bar{U} \subset V_x \subset \bar{V}_x \subset U'$ , for which the difference of estimates of  $\alpha(V_x, x, p)$  in

$$\alpha(U', x, p) \leq \alpha(V_x, x, p) \leq \alpha(U, x, p)$$

is arbitrarily small. For  $\delta(V_x, V_y)$  small enough we have  $U \subset V_y \subset U'$  and the analogous estimate together with the continuity of  $\omega$  gives

$$(13) \quad |\alpha(V_x, x, p) - \alpha(V_y, x, p)| \rightarrow 0$$

as  $\rho(x, y) \rightarrow 0$ . It is easily seen that for all  $y$  sufficiently close to  $x$  we have  $U \subset V_y$  and hence the functions  $\alpha(V_y, \cdot, p)$  are harmonic on  $U$ . Moreover, these functions are uniformly bounded for all  $y$  from a neighbourhood of  $x$ . This implies by Th. 11.1.1 in [3] the equicontinuity of all these functions on a neighbourhood of  $x$ . Now from (13), the equicontinuity and the estimate

$$|Ap(x) - Ap(y)| \leq |\alpha(V_x, x, p) - \alpha(V_y, x, p)| + |\alpha(V_y, x, p) - \alpha(V_y, y, p)|$$

we shall easily obtain the required continuity of  $Ap$  at  $x$ .

In the case of classical harmonic functions the mean value over a ball can be interpreted as an "infinite combination" of harmonic measures corresponding to smaller balls with the same centre. In what follows, we shall study  $A$ -harmonicity for  $A = \{\alpha_x\}_{x \in G}$  with measures  $\alpha_x$  of a similar type.

8. Lemma. Let  $D$  be an open subset of a metric space. Let us suppose:

- (i)  $\{f_x; x \in D\}$  is a set of positive decreasing and uni-

formly bounded functions defined on  $[0, d]$ ,  $d > 0$ ;

(ii) for any  $x \in D$  and  $s, t, t' \in [0, d]$ ,  $t < s < t'$ , there is a neighbourhood  $U(x)$  of  $x$  in  $D$  such that the inequality

$$(14) \quad f_x(t) \geq f_y(t) \geq f_x(t')$$

holds for every  $y \in U(x)$ ;

(iii)  $g$  is a continuous strictly increasing function on  $[0, d]$ ,  $g(0) = 0$ .

Then the average

$$(15) \quad \mathcal{A}(f_x, g, c) := (g(c))^{-1} \int_0^c f_x(t) dg(t)$$

is a continuous function (w. r. t. the usual product topology) of  $(x, c)$  on  $D \times ]0, d[$ .

Proof. Clearly,  $c \mapsto \mathcal{A}(f_x, g, c)$  is a continuous function on  $]0, d[$  for any  $x \in D$ . It is easily seen that it is also decreasing. For a fixed  $c \in ]0, d[$  the continuity of  $x \mapsto \mathcal{A}(f_x, g, c)$  follows from the approximation of the integral in (15) by an integral sum combined with the property (14). Indeed, for  $P \equiv \{0 = t_0 < t_1 < \dots < t_k = c\}$  and  $s_1 \in ]t_{i-1}, t_i[$  we can choose a neighbourhood  $U(x)$  of  $x$  such that  $f_x(t_{i-1}) \geq f_y(s_1) \geq f_x(t_i)$  holds for all  $y \in U(x)$ . Now the expression

$$\left| \sum_{i=1}^k f_x(t_i) [g(t_i) - g(t_{i-1})] - \sum_{i=1}^k f_y(s_i) [g(t_i) - g(t_{i-1})] \right|$$

may be done arbitrarily small by the choice of  $P$  and  $s_i$  while both integral sums are close enough to the corresponding integrals. (For details see the proof of L.11 in [13].) From the separate continuity in  $x$  and  $c$  and monotonicity in  $c$  it follows that  $\mathcal{A}(f_x, g, c)$  is continuous on  $D \times ]0, d[$ .

9. Proposition. Let  $G \subset X$  be an open set. Let us suppose that

$$\Omega: G \times G \rightarrow \mathbb{R}^1, \quad d: G \rightarrow \mathbb{R}^1, \quad g: [0, \infty[ \rightarrow \mathbb{R}^1$$

are positive continuous functions enjoying properties:

(i)  $\Omega(x, x) = 0$  for every  $x \in G$ ,  $\Omega(x, y) > 0$  for every  $x, y \in G$ ,  $x \neq y$ ;

(ii) for every  $x \in G$ ,  $0 < c < d(x)$ , the closure of the set

$$(16) \quad V_{x,c} = \{y \in G; \Omega(x, y) < c\}$$

is a compact subset of  $G$  and its distance (w.r.t. the metric  $\rho$  fixed above) from  $\partial V_{x,c}$ , is strictly positive for every  $c < c' < d(x)$ ;

(iii)  $g$  is a continuous strictly increasing function and  $g(0) = 0$

Then for every  $h \in \mathcal{C}(G)$

$$(17) \quad Ah(x, c) := (g(c))^{-1} \int_0^c \infty(V_{x,t}, x, h) dg(t)$$

defines a continuous function of  $(x, c)$  on the set  $\{(x, c); x \in G, 0 < c < d(x)\}$ .

*Proof.* Fix a point  $(x, c)$  in the set and choose a neighbourhood  $D$  of  $x$  and an interval  $]a, b[$  containing  $c$  in such a way that  $\bar{V}_{x,e} \subset G$  for all  $(y, e) \in D \times ]a, b[$ .

Likewise in the proof of Pr. 7 it suffices to prove the continuity of  $A_p$  for every potential  $p$ . We shall show that in this case Pr. 9 follows easily from L.8. Indeed, the assumption (iii) implies that L.8(iii) holds for  $g$  on  $[0, b]$ . On the same interval, the functions  $f_x: t \mapsto \infty(V_{x,t}, x, p)$ ,  $x \in D$ , satisfy L.8(i).

If  $x \in D$  and  $s, t, t' \in [0, b]$ ,  $t < s < t'$ , then it follows from (ii) and the uniform continuity of  $\Omega$  on compact subsets of  $G \times G$  that there is a neighbourhood  $U(x)$  of  $x$  in  $D$  such that we have  $V_{x,t} \subset V_{y,s} \subset V_{x,t'}$  for every  $y \in U(x)$ . Now (14) is a consequence of the monotonicity of the balayage. Since we have verified that all assumptions of L.8 are fulfilled, the proposition is proved.

10. Corollary. Assume the same as in Pr. 9 and suppose that  $\sigma \in \mathcal{B}(G)$ ,  $0 < \sigma(x) < d(x)$  for every  $x \in G$ . Then

$$(18) \quad A_1 h(x) := Qh(x, \sigma(x)), \quad x \in G,$$

where  $Qh$  is defined by (17), and

$$(19) \quad A_2 h(x) := \alpha(V_{x, \sigma(x)}, x, h), \quad x \in G$$

are Borel operators (on  $\mathcal{B}_b(G)$ ). If  $\sigma$  is moreover continuous, then  $A_1 h \in \mathcal{C}_b(G)$  and  $A_2 h \in \mathcal{B}_b(G)$  whenever  $h \in \mathcal{C}_b(G)$ .

Indeed, if  $h \in \mathcal{C}_b(G)$  and  $\sigma$  is continuous, then Pr. 9 implies that  $Qh(x, \sigma(x))$ ,  $x \in G$  is continuous. Put  $g(t) = t$ ,  $t > 0$ . It is easily seen that

$$A_2 h(x) = \lim_{n \rightarrow \infty} n (Qh(x, \sigma(x)) - Qh(x, \sigma(x) - n^{-1}))$$

and hence  $A_2 h \in \mathcal{B}_b(G)$ . This gives the first part of the assertion.

Now we can give a more explicit description of a broad class of operators  $A$  which we met in connection with  $A$ -harmonicity in Th. 6. More precisely, we shall show how to construct for the operators  $A_1, A_2$  from (18) and (19) a corresponding  $L$ -operator  $B$  for which (9) holds provided "the values of the size function  $\sigma$  are not too small" in  $G$ . The same will be shown

for an operator A from Pr. 7.

11. Theorem. Let  $G \subset X$  be an open relatively compact set and  $\rho$  the metric fixed above.

(a) Assume, moreover, the same on  $\Omega$ ,  $d$ ,  $g$  and  $\sigma$  as what was supposed in Pr. 9 and Cor. 10. If for every compact set  $C \subset G$

$$(20) \quad \inf \{ \sigma(x); x \in C \} = i_C > 0,$$

then the operators  $A_1, A_2$  defined by (18) and (19) have all properties required in Th. 6.

(b) Assume now the same on  $\omega$  and  $V_x$  what was supposed in Pr. 7. If

$$(21) \quad \sigma(x) = \text{dist}(x, CV_x)$$

fulfils the condition (20), then the operator A from Pr. 7 has all properties required in Th. 6.

Proof. All what remains to prove is the existence of a corresponding operator B. In the first case (i.e. (a)) choose compact sets  $C_k, C_k \nearrow G$  and for  $k = 1, 2, \dots$  put  $i_k := (k+1)^{-1} i_{C_k}$  (cf. (20)). Then choose  $f_k \in \mathcal{C}_0(X)$  in such a way that  $0 \leq f_k \leq i_k - i_{k+1}$ ,  $f_k(x) = i_k - i_{k+1}$  on  $C_k$  and  $f_k(x) = 0$  on  $CC_{k+1}$ . Now denote by  $\Delta$  the restriction of  $\sum f_k$  on  $G$ ; since  $\Delta \in \mathcal{C}(G)$ ,  $0 < \Delta(x) < \sigma(x)$ , B can be defined by

$$Bh(x) := \mathcal{A}h(x, \Delta(x)), \quad x \in G$$

(and, formally,  $Bh(x) = h(x)$ ,  $x \in \partial G$ ) and every  $h \in \mathcal{C}(\bar{G})$ . Inequality (9) is a consequence of monotonicity of  $\mathcal{A}h(x, c)$  from (17) in the variable  $c$ .

In case of (b) the construction can be repeated with  $\Omega(x,y) = \varphi(x,y)$ ,  $d(x) = \text{dist}(x,CG)$  and  $g(t) = t$ ,  $t > 0$ . Function  $d'$  is defined by (21). Inequality (9) now follows from the monotonicity of the balayage.

#### 12. Remarks and comments

(1) Lebesgue in [7] replaced essentially the Laplace equation in the Dirichlet problem by the equation  $Af = f$  with a "Lebesgue's operator"  $A$ . He obtained one of the first theorems on the restricted mean value property (cf. also [8]).

(2) Our results are a generalization of a theorem proved in the classical case in [11]; the authors use a probabilistic approach. Remark that in  $R^n$  the mean values of  $h$  w.r.t. balls can be obtained if we choose  $\Omega(x,y) = \varphi(x,y)$ , the Euclidean metric and  $g(t) = t^{n-1}$ ,  $t > 0$ ; then (17) determines the mean value since in the case  $\omega(V_{x,t}, \cdot)$  is the normalized surface measure on the sphere  $\partial V_{x,t}$ .

(3) For the generalization of the Lebesgue's approach to the Dirichlet problem in [7] see [13]. In the formula (20) in [13] instead of  $\varepsilon_x^{CD} t$  should be  $\varepsilon_x^{CD}(t)$ .

(4) In our assertions we used mean values of a quite general form but we admit only functions with a "good boundary behaviour a.e. on  $\partial G$ ". It would be interesting to know whether similar results can be obtained in the frame of an axiomatics for functions with more general boundary behaviour.

#### R e f e r e n c e s

- [1] ASH R.B.: Measure, Integration and Functional Analysis, Academic Press, New York and London 1972.



- [2] BAUER H.: *Harmonische Räume und ihre Potentialtheorie*, Springer Verlag, Berlin 1966.
- [3] CONSTANTINESCU C., CORNE A.: *Potential Theory on Harmonic Spaces*, Springer Verlag, New York 1972.
- [4] FENTON P.C.: On sufficient conditions for harmonicity, *Trans. Amer. Math. Soc.* 253(1979), 139-147.
- [5] HEATH D.: Functions possessing restricted mean value properties, *Proc. Amer. Math. Soc.* 41(1973), 588-595.
- [6] KELLOG O.D.: Converses of Gauss's theorem on the arithmetic mean, *Trans. Amer. Math. Soc.* 36(1934), 227-242.
- [7] LEBESGUE H.: Sur le problème de Dirichlet, *C.R. Acad. Sci. Paris* 154(1912), 335-337.
- [8] LEBESGUE H.: Sur le théorème de la moyenne de Gauss, *Bull. Soc. Math. France* 40(1912), 16-17.
- [9] NETUKA I.: Harmonic functions and the mean value theorems (in Czech), *Čas. pěst. mat.* 100(1975), 391-409.
- [10] NETUKA I.: L'unicité du problème de Dirichlet généralisé pour un compact, in: *Séminaire de Théorie du Potentiel Paris*, No. 6, *Lecture Notes in Mathematics* 906, Springer Verlag, Berlin 1982, 269-281.
- [11] ØKSENDAL B., STROOCK D.W.: A characterization of harmonic measure and Markov processes whose hitting distributions are preserved by rotations, translations and dilatations (preprint).
- [12] VEECH W.A.: A converse to the mean value theorem for harmonic functions, *Amer. J. Math.* 97(1976), 1007-1027.
- [13] VESELÝ J.: Sequence solutions of the Dirichlet problem, *Čas. pěst. mat.* 106(1981), 84-93.

Matematicko-fyzikální fakulta UK, Sokolovská 83, 18600 Praha 8, Czechoslovakia

(Oblatum 25.8. 1982)