

## Werk

**Label:** Article

**Jahr:** 1987

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?316342866\\_0028|log33](https://resolver.sub.uni-goettingen.de/purl?316342866_0028|log33)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

AROUND A NEUTRAL ELEMENT IN A NEARLATTICE  
A.S.A. NOOR and W.H. CORNISH

Abstract: Nearlattices, or lower semilattices in which any two elements have a supremum whenever they are bounded above, provide an interesting generalization of lattices. In this context, we define different types of elements in a nearlattice  $S$  and then for a fixed element  $n$ , using the ternary operation  $J_n$ , study the behaviour of  $S_n = (S; \cap)$  where  $x \cap y = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ ;  $x, y \in S$ .

Key words: Standard element, neutral element, nearlattice.  
Classification: 06A12, 06A99, 06B10

1. Introduction. A nearlattice is a lower semilattice which has the property that any two elements possessing a common upper bound, have a supremum. Cornish and Hickman [1] called this the upper bound property. For detailed literature, we refer the reader to consult [1], [2] and [7].

A nearlattice-congruence  $\Phi$  on a nearlattice  $S$  is a congruence of the underlying lower semilattice such that, whenever  $a_1 \equiv b_1, a_2 \equiv b_2 (\Phi)$  and  $a_1 \vee a_2, b_1 \vee b_2$  exist,  $a_1 \vee a_2 \equiv b_1 \vee b_2 (\Phi)$ . In the second section of [4], a fundamental contribution was made by Hickman. Defining a ternary operation  $j$  on a nearlattice  $S$  by  $j(x, y, z) = (x \wedge y) \vee (y \wedge z)$ , he showed that the resulting algebras of the type  $(S; j)$  form a variety.

Standard and neutral elements, as well as standard ideals in a nearlattice were extensively studied in [2]. An element  $s$  in a nearlattice  $S$  is called standard if for all  $x, y, t \in S$ ,  $t \wedge [(x \wedge y) \vee (x \wedge s)] = (t \wedge x \wedge y) \vee (t \wedge x \wedge s)$ . An element  $n$  in a nearlattice  $S$  is called neutral if it is standard and for any  $t, x, y \in S$ ,  $n \wedge [(t \wedge x) \vee (t \wedge y)] = (n \wedge t \wedge x) \vee (n \wedge t \wedge y)$ . Clearly, every element of a distributive nearlattice is neutral. An ele-

ment  $n$  of a nearlattice  $S$  is called superstandard if it is standard and for any  $x, y \in S$ ,  $n \wedge [(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)] = (x \wedge n) \vee (y \wedge n)$ , whenever  $(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$  exists. Of course, every neutral element is superstandard. But in the pentagonal lattice  $\{0, a, b, n, 1\}$  where  $0 < a < n < 1$ ;  $0 < b < 1$ :  $a \wedge b = n \wedge b = 0$  and  $a \vee b = n \vee b = 1$ ,  $n$  is superstandard but not neutral. [7] provides an example of a standard element in a lattice which is not superstandard.

An element  $n$  in a nearlattice  $S$  is called medial if  $m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$  exists for all  $x, y \in S$ , while  $n$  is called sesquimedial if  $J_n(x, y, z) = [(x \wedge n) \vee (y \wedge n)] \wedge [(y \wedge n) \vee (z \wedge n)] \vee j(x, y, z)$  exists for all  $x, y, z \in S$  where  $j(x, y, z) = (x \wedge y) \vee (y \wedge z)$ . Since  $J_n(x, y, x) = m(x, n, y)$  for all  $x, y \in S$ , any sesquimedial element is medial. A nearlattice  $S$  is called medial if  $m(x, y, z) = (x \wedge y) \vee (y \wedge z) \vee (z \wedge x)$  exists for all  $x, y, z \in S$ . Of course, every element of a medial nearlattice is sesquimedial (see Lemma 3.1).

Let  $n$  be a fixed element of a nearlattice  $S$ . By an  $n$ -ideal of  $S$ , we mean a convex subnearlattice of  $S$  containing  $n$ . The  $n$ -ideal generated by  $a_1, \dots, a_m$  is denoted by  $\langle a_1, \dots, a_m \rangle_n$ . Clearly  $\langle a_1, \dots, a_m \rangle_n = \langle a_1 \rangle_n \vee \dots \vee \langle a_m \rangle_n$ . When  $S$  is a lattice,  $\langle a_1, \dots, a_m \rangle_n = \langle a_1 \wedge \dots \wedge a_m \wedge n, a_1 \vee \dots \vee a_m \vee n \rangle_n$ . Thus, for a lattice  $S$ , the set of finitely generated  $n$ -ideals of  $S$  is a lattice and its members are simply the intervals  $[a, b]$  such that  $a \leq n \leq b$ , and for such intervals,  $[a, b] \vee [a_1, b_1] = [a \wedge a_1, b \vee b_1]$  and  $[a, b] \cap [a_1, b_1] = [a \vee a_1, b \wedge b_1]$ . The  $n$ -ideal generated by a single element is called a Principal  $n$ -ideal and the set of Principal  $n$ -ideals of  $S$  is denoted by  $P_n(S)$ . When  $S$  is a lattice, it is not hard to see that  $P_n(S)$  is a lattice if and only if  $n$  is complemented in each interval containing it.

For a fixed element  $n$ , the binary operation  $x \circ y = m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$  has been studied by several authors including Jakubík and Kolibiar [5] for distributive lattices, Sholander [8] for distributive medial near lattices and Kolibiar [6] for an arbitrary lattice with  $n$  as a neutral element in it. Sholander [8] showed that for a distributive medial nearlattice  $S$ ,  $(S; \circ)$  is a semilattice. On the other hand Kolibiar [6] showed that if  $n$  is a neutral element in an arbitrary lattice  $S$ ,  $(S; \circ)$  is a semilattice. Recently, Noor [7] extended their work and showed that for a neutral and sesquimedial element  $n$  of a near-

lattice  $S$ ,  $S_n = (S; \cap)$  is not only a semilattice, it is a nearlattice. Moreover, the  $n$ -ideals of  $S$  are precisely the ideals of  $S_n$ . According to [7], we refer to  $S_n$  as an isotope of  $S$ .

In Section 2, we introduce the notion of a nearly neutral element in a nearlattice and then generalize and extend some of the results in [7]. We show that for a medial superstandard element  $n$  of a nearlattice  $S$ ,  $S_n$  is a nearlattice wherein  $J_n(x, y, z) = j_n(x, y, z)$  if and only if  $n$  is nearly neutral and sesquimedial in  $S$ . We also show that for a nearly neutral and sesquimedial element of a nearlattice  $S$ ,  $n$  is neutral if and only if the nearlattice congruences of  $S$  are precisely the nearlattice congruences of  $S_n$ .

In Section 3, introducing the ternary operation  $M_n(x, y, z)$  we show that for a sesquimedial neutral element  $n$  of a nearlattice  $S$ ,  $S$  is medial if and only if  $S_n$  is so.

**2. Nearly neutral element of a near lattice.** An element  $n$  of a nearlattice is called nearly neutral if it is standard and has the property  $n \wedge ((t \wedge x \wedge n) \vee (t \wedge y)) = (t \wedge x \wedge n) \vee (t \wedge y \wedge n)$  for all  $x, y, t \in S$ . Of course, a neutral element is always nearly neutral. Observe that in Figure 1,  $n$  is nearly neutral but  $n \wedge (a \vee b) > (n \wedge a) \vee (n \wedge b)$  shows that it is not neutral there.

The following result shows that every nearly neutral element is superstandard, but in the pentagonal lattice  $\{0, a, b, n, 1\}$  where  $0 < a < n < 1$ ;  $0 < b < 1$ ;  $a \wedge b = n \wedge b = 0$ ;  $a \vee b = n \vee b = 1$ ,  $n$  is superstandard but not nearly neutral.

**Proposition 2.1.** For an element  $n$  of a nearlattice  $S$ , the following conditions are equivalent.

- (i) For all  $x, y, t \in S$ ,  
 $n \wedge ((t \wedge x \wedge n) \vee (t \wedge y)) = (t \wedge x \wedge n) \vee (t \wedge y \wedge n)$ .
- (ii) For all  $x, y \in S$ ,  
 $n \wedge ((x \wedge n) \vee y) = (x \wedge n) \vee (y \wedge n)$ , whenever  $(x \wedge n) \vee y$  exists.

Moreover, if  $n$  is sesquimedial, (i) and (ii) are also equivalent to each of the next two conditions.

- (iii) For all  $x, y, z \in S$ ,  $(x \cap y) \wedge n = (x \wedge n) \vee (y \wedge n)$  and  $J_n(x, y, z) \wedge n = (x \cap y) \wedge (y \cap z) \wedge n$ , where  $x \cap y = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ .

(iv) For all  $x, y, z \in S$ ,  $(x \circ y) \wedge n = (x \wedge n) \vee (y \wedge n)$  and  $J_n(x, y, z) \wedge n \leq x \circ y$ .

Proof. (i)  $\Rightarrow$  (ii). Suppose  $(x \wedge n) \vee y$  exists. Then  $n \wedge ((x \wedge n) \vee y) = n \wedge [((x \wedge n) \vee y) \wedge x \wedge n] \vee [((x \wedge n) \vee y) \wedge y] = (x \wedge n) \vee (y \wedge n)$ .  
(ii)  $\Rightarrow$  (i) is trivial.

Suppose now that  $n$  is sesquimedial and (i) and (ii) hold. Then  $n \wedge (x \circ y) = n \wedge ((x \wedge n) \vee (y \wedge n) \vee (x \wedge y)) = n \wedge [((x \wedge n) \vee (y \wedge n)) \wedge n] \vee (x \wedge y) = (x \wedge n) \vee (y \wedge n) \vee (x \wedge y \wedge n) = (x \wedge n) \vee (y \wedge n)$ . Also,  $J_n(x, y, z) \wedge n = n \wedge [((x \wedge n) \vee (y \wedge n)) \wedge ((y \wedge n) \vee (z \wedge n))] \vee (x \wedge y) \vee (y \wedge z) = n \wedge [((x \circ y) \wedge (y \circ z)) \wedge n] \vee (x \wedge y) \vee (y \wedge z) = ((x \circ y) \wedge (y \circ z)) \vee (n \wedge ((x \wedge y) \vee (y \wedge z))) = (x \circ y) \wedge (y \circ z) \wedge n$ .

Thus (iii) holds.

Clearly (iii) implies (iv).

Finally suppose (iv) holds. Let  $x, y \in S$  be such that  $(x \wedge n) \vee y$  exists. Then

$J_n(x, y, (x \wedge n) \vee y) = [((x \wedge n) \vee (y \wedge n)) \wedge (y \wedge n) \vee (n \wedge ((x \wedge n) \vee y))] \vee (x \wedge y) \vee y = (x \wedge n) \vee (y \wedge n) \vee y = (x \wedge n) \vee y$ , and so by (iv)  $n \wedge ((x \wedge n) \vee y) \leq x \circ y$ . Thus,  $n \wedge ((x \wedge n) \vee y) \leq n \wedge (x \circ y) = (x \wedge n) \vee (y \wedge n)$ ; it follows that  $n \wedge ((x \wedge n) \vee y) = (x \wedge n) \vee (y \wedge n)$  and (ii) holds.  $\square$

The following result is found in [7, Th. 2.1].

Proposition 2.2. If  $n$  is a standard element of a nearlattice  $S$ , then  $(S; \leq)$  is a partially ordered set and the map  $x \rightarrow \langle x \rangle_n$  is an isomorphism of  $(S; \leq)$  onto  $P_n(S)$ , where on  $S$ ,  $x \leq y$  if and only if  $(x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$  exists and is equal to  $x$ .  $\square$

Let  $n$  be a medial element of a nearlattice  $S$ . For any  $x, y \in S$  define the binary operation  $x \circ y = m(x, n, y) = (x \wedge y) \vee (x \wedge n) \vee (y \wedge n)$ . Recently Noor in [7] proved the following result.

Theorem 2.3. If  $n$  is a medial and standard element of a nearlattice  $S$ , then  $S_n$  is a semilattice if and only if  $n$  is superstandard in  $S$ .

Moreover, when  $n$  is neutral and sesquimedial then  $S_n$  is in fact a nearlattice and the  $n$ -ideals of  $S$  are precisely the ideals of  $S_n$ .  $\square$

Our next theorem generalizes and extends the above Theorem. To obtain this, we need the following lemma. (i) is found in

[7; Lemma 2.4], and the proof of (ii) is similar to the proof of (i) in [7; Lemma 2.4].

Lemma 2.4. In a nearlattice  $S$ ,

(i) a subset  $K$  of  $S$  is an ideal of  $S$  if and only if for all  $x, y \in K$  and  $a \in S$ ,  $j(x, a, y) \in K$ .

(ii) If  $n$  is a superstandard element of  $S$  such that  $S_n$  is a nearlattice wherein

$$J_n(x, y, z) = j^{S_n}(x, y, z) = (x \cap y) \cup (y \cap z),$$

then a subset  $K$  of  $S$  is an  $n$ -ideal of  $S$  if and only if it contains  $n$  and  $J_n(x, a, y) \in K$  for any  $x, y \in K$  and  $a \in S$ .  $\square$

Corollary 2.5. Suppose  $n$  is a superstandard element of a nearlattice  $S$  such that the isotope  $S_n$  of  $S$  is itself a nearlattice wherein  $J_n(x, y, z) = j^{S_n}(x, y, z)$ . Then the ideals of  $S_n$  are precisely the  $n$ -ideals of  $S$ .  $\square$

Theorem 2.6. If  $S$  is a nearlattice and  $n \in S$  is medial and superstandard, then the following conditions are equivalent.

(i)  $n$  is nearly neutral and sesquimedial in  $S$ .

(ii) The isotope  $S_n = (S; \cap)$  is a nearlattice wherein

$$J^{S_n}(x, y, z) = J_n(x, y, z).$$

(iii)  $S_n$  has the upper bound property and  $n$ -ideals of  $S$  are precisely the ideals of  $S_n$ .

(iv) Any finitely generated  $n$ -ideal contained in a principal  $n$ -ideal is a principal  $n$ -ideal.

Proof. (i)  $\Rightarrow$  (ii). Suppose  $n$  is nearly neutral and sesquimedial in  $S$ . Then, clearly

$$J_n(x, y, z) = ((x \cap y) \cap (y \cap z) \cap n) \vee j(x, y, z),$$

and so by [2; Th. 2.4],  $J_n(x, y, z) \equiv j(x, y, z)(\Theta_n)$  and

$x \cap y \equiv x \wedge y(\Theta_n)$ . Hence  $(x \cap y) \wedge J_n(x, y, z) \equiv x \wedge y(\Theta_n)$  and simi-

larly  $(y \cap z) \wedge J_n(x, y, z) \equiv y \wedge z(\Theta_n)$ . Therefore,

$$\begin{aligned} [(x \cap y) \wedge J_n(x, y, z)] \vee [(y \cap z) \wedge J_n(x, y, z)] \vee [n \wedge J_n(x, y, z)] &\equiv \\ \equiv (x \wedge y) \vee (y \wedge z) \vee (n \wedge j(x, y, z)) &= j(x, y, z)(\Theta_n). \end{aligned}$$

Since the left hand side of this congruence exceeds the right hand side, by [2; Th. 2.4],

$$\begin{aligned} &\text{left hand expression} \\ &= j(x, y, z) \vee (n \wedge (\text{left hand expression})) \end{aligned}$$

$$= j(x,y,z) \vee (n \wedge J_n(x,y,z)) = J_n(x,y,z).$$

Thus,  $J_n(x,y,z) \in \langle x\wedge y, y\wedge z \rangle_n$ . On the other hand,  $(x\wedge y) \wedge J_n(x,y,z) \equiv x\wedge y(\Theta_n)$  implies  $(x\wedge y) \wedge J_n(x,y,z) = (x\wedge y) \vee (n \wedge (x\wedge y) \wedge J_n(x,y,z))$ , and so  $((x\wedge y) \wedge J_n(x,y,z)) \vee ((x\wedge y) \wedge n) = (x\wedge y) \vee (x \wedge n) \vee (y \wedge n) = x\wedge y$ . Hence,  $x\wedge y \in \langle J_n(x,y,z) \rangle_n$  and similarly  $y\wedge z \in \langle J_n(x,y,z) \rangle_n$ . Thus,  $\langle x\wedge y, y\wedge z \rangle_n = \langle J_n(x,y,z) \rangle_n$  and so by Proposition 2.2,  $(x\wedge y) \vee (y\wedge z) = J_n(x,y,z)$ .

(ii)  $\Rightarrow$  (iii) follows immediately from Corollary 2.5.

(iii)  $\Rightarrow$  (iv) is an easy consequence of the isomorphism of  $(S_n; \subseteq)$  and  $(P_n(S); \subseteq)$ , and the upperbound property of  $S_n$ .

(iv)  $\Rightarrow$  (i). Let  $a, b, c \in S$ . Since  $a \wedge b, b \wedge c \in b$ ,  $\langle a \wedge b, b \wedge c \rangle_n \subseteq \langle b \rangle_n$  by Proposition 2.2. Thus, by (iv), there exists  $t \in S$  such that  $\langle a \wedge b, b \wedge c \rangle_n = \langle t \rangle_n$ , and so  $(a \wedge b) \wedge (b \wedge c) \wedge n = t \wedge n$ . Now,  $a \wedge b \leq t$  implies  $a \wedge b = ((a \wedge b) \wedge t) \vee (a \wedge b) \wedge n = ((a \wedge b) \wedge t) \vee ((a \wedge b) \wedge n)$ , and so  $a \wedge b \equiv (a \wedge b) \wedge t(\Theta_n)$ . Hence  $a \wedge b \equiv a \wedge b \equiv (a \wedge b) \wedge t \equiv a \wedge b \wedge t(\Theta_n)$ .

Similarly,  $b \wedge c \equiv b \wedge c \wedge t(\Theta_n)$ . This implies

$$j(a,b,c) \equiv (a \wedge b \wedge t) \vee (b \wedge c \wedge t) (\Theta_n)$$

and so  $j(a,b,c) = (a \wedge b \wedge t) \vee (b \wedge c \wedge t) \vee (n \wedge j(a,b,c))$ . Also,  $j(a,b,c) \wedge t \equiv (a \wedge b \wedge t) \vee (b \wedge c \wedge t) (\Theta_n)$ , and so

$$j(a,b,c) \wedge t = (a \wedge b \wedge t) \vee (b \wedge c \wedge t) \vee (n \wedge t \wedge j(a,b,c)).$$

Thus,  $j(a,b,c) \wedge t = (j(a,b,c) \wedge t) \vee (j(a,b,c) \wedge n) \vee (t \wedge n) = j(a,b,c) \vee (t \wedge n)$ .

Again,  $a \wedge b \equiv a \wedge b(\Theta_n)$ . So  $(a \wedge b) \wedge j(a,b,c) \equiv a \wedge b \wedge j(a,b,c) = a \wedge b(\Theta_n)$ , and hence  $(a \wedge b) \wedge j(a,b,c) = (a \wedge b) \vee ((a \wedge b) \wedge j(a,b,c) \wedge n)$ . This implies  $(a \wedge b) \wedge j(a,b,c) = a \wedge b$ ; that is,  $a \wedge b \leq j(a,b,c)$ . Similarly,  $b \wedge c \leq j(a,b,c)$ . Hence,  $t \leq j(a,b,c)$ , and so  $t = t \wedge j(a,b,c) = j(a,b,c) \vee (t \wedge n) = j(a,b,c) \vee ((a \wedge b) \wedge (b \wedge c) \wedge n) = J_n(a,b,c)$ , as  $n$  is superstandard. Hence  $n$  is sesquimedial, and  $J_n(a,b,c) \wedge n = t \wedge n = (a \wedge b) \wedge (b \wedge c) \wedge n$ . Also  $(x \wedge y) \wedge n = (x \wedge n) \vee (y \wedge n)$ , as  $n$  is superstandard. Therefore, by 2.1(iii),  $n$  is nearly neutral.  $\square$

The following lemma is due to Hickman [4; Proposition 2.2].

**Lemma 2.7.** In a nearlattice  $S$ , an equivalence relation is a nearlattice congruence if and only if it is a congruence for the algebra  $(S; j)$ .  $\square$

Now we consider the influence of  $J_n$  on congruences. The following theorem is an extension of [7; Lemma 2.6(ii)].

**Theorem 2.8.** Let  $n$  be a sesquimedial, nearly neutral element of a nearlattice  $S$ . Then the following conditions are equivalent.

- (i)  $n$  is neutral in  $S$ ;
- (ii) an equivalence relation on  $S$  is a congruence for the algebra  $(S; J_n)$  if and only if it is a nearlattice-congruence of  $S$ .

**Proof.** (i)  $\Rightarrow$  (ii) is proved in [7; Lemma 2.6(ii)].

(ii)  $\Rightarrow$  (i). Define a relation  $\Theta$  on the nearlattice  $S$  by  $x \equiv y(\Theta)$  if and only if  $x \wedge n = y \wedge n$ . This is clearly an equivalence relation on  $S$ .

Now suppose  $x \equiv y(\Theta)$ . Then  $x \wedge n = y \wedge n$ , and so by 2.1, for any  $s, t \in S$ ,  $n \wedge J_n(x, s, t) = (x \wedge n) \wedge (s \wedge t) \wedge n = ((x \wedge n) \vee (s \wedge n)) \wedge ((s \wedge n) \vee (t \wedge n)) = ((y \wedge n) \vee (s \wedge n)) \wedge ((s \wedge n) \vee (t \wedge n)) = n \wedge J_n(y, s, t)$ . Thus,  $J_n(x, s, t) \equiv J_n(y, s, t)(\Theta)$ . Similarly,  $J_n(s, x, t) \equiv J_n(s, y, t)(\Theta)$  and  $J_n(s, t, x) \equiv J_n(s, t, y)(\Theta)$ , and so  $\Theta$  is a congruence for the algebra  $(S; J_n)$ . Thus, by (ii),  $\Theta$  is a nearlattice congruence on  $S$ . Now, clearly  $x \equiv x \wedge n(\Theta)$  and  $y \equiv y \wedge n(\Theta)$  for all  $x, y \in S$ . So for any  $t \in S$ ,  $(t \wedge x) \vee (t \wedge y) \equiv (t \wedge x \wedge n) \vee (t \wedge y \wedge n)(\Theta)$ , and hence,  $n \wedge [(t \wedge x) \vee (t \wedge y)] = n \wedge [(t \wedge x \wedge n) \vee (t \wedge y \wedge n)] = (t \wedge x \wedge n) \vee (t \wedge y \wedge n)$ , which implies  $n$  is neutral in  $S$ .  $\square$

Combining Theorem 2.6, Lemma 2.7 and the above theorem, we have the following extension of [7, Th. 2.7].

**Theorem 2.9.** Let  $n$  be a nearly neutral sesquimedial element of a nearlattice  $S$ . Then  $n$  is neutral if and only if the nearlattice congruences of  $S$  are precisely the nearlattice congruences of  $S_n$ .  $\square$

The following proposition will be needed to prove one of our main results in Section 3. This was known by Kolibiar [6] in case of a bounded lattice with  $n$  as a central element.

**Proposition 2.10.** If  $n$  is a nearly neutral sesquimedial element of a nearlattice  $S$  with  $0$ , then  $0$  is neutral and medial in  $S_n$ . Moreover, the double isotope  $(S_n)_0$  is precisely  $S$ .

If, in addition,  $n$  is neutral in  $S$ , then  $0$  is sesquimedial in  $S_n$  and  $J_0^{S_n}(x, y, z) = j_0^{(S_n)}(x, y, z) = j(x, y, z) = J_0(x, y, z)$  for all  $x, y, z \in S$ .



Proof. By 2.6, for all  $r, x, y \in S$ ,  $0 \cap ((r \cap x) \cup (r \cap y)) =$   
 $= 0 \cap J_n(x, r, y) = n \wedge J_n(x, r, y) = J_n(r \cap x, 0, r \cap y) = (0 \cap r \cap x) \cup (0 \cap r \cap y)$ .  
 Also,  $r \cap ((x \cap y) \cup (x \cap 0)) = r \cap J_n(y, x, 0) = r \cap ((x \cap n) \vee (x \cap y)) = (r \cap n) \vee$   
 $\vee (x \cap n) \vee (r \cap x \cap y)$  as  $n$  is nearly neutral and hence standard. On the  
 other hand,  $(r \cap x \cap y) \cup (r \cap x \cap 0) = J_n(y, r \cap x, 0) = ((r \cap x) \wedge n) \vee (y \wedge (r \cap x)) =$   
 $= (r \cap n) \vee (x \cap n) \vee [y \wedge ((r \cap x) \wedge r \cap x) \vee ((r \cap x) \wedge n)] = (r \cap n) \vee (x \cap n) \vee (r \cap x \cap y)$ .  
 That is  $r \cap ((x \cap y) \cup (x \cap 0)) = (r \cap x \cap y) \cup (r \cap x \cap 0)$ ; consequently  $0$  is ne-  
 utral in  $S_n$ .

Now, clearly  $x \cap y, x \cap 0, y \cap 0 \in x \cap y$ , and so  $(x \cap y) \cup (x \cap 0) \cup (y \cap 0)$  ex-  
 ists and it is  $\in x \cap y$ . Thus,  $0$  is medial in  $S_n$ , and so  $((S_n)_0; \bar{\wedge})$   
 is a semilattice by Theorem 2.3, where

$$x \bar{\wedge} y = (x \cap y) \cup (x \cap 0) \cup (y \cap 0).$$

Suppose  $x \cap y, x \cap 0, y \cap 0 \in s$  for some  $s \in S_n$ . Then  $s \wedge n \in (x \cap 0) \wedge n =$   
 $= x \cap n$ . Similarly  $s \wedge n \in y \cap n$ , and so  $s \wedge n \in x \cap y \cap n$ . Also,  
 $x \cap y = (x \cap y) \cap s = ((x \cap y) \wedge s) \vee ((x \cap y) \wedge n) \vee (s \wedge n) =$   
 $= ((x \cap y) \wedge s) \vee ((x \cap y) \wedge n)$ .

Then

$$x \cap y = (x \cap y) \wedge (x \cap y) = (x \cap y \wedge s) \vee (x \cap y \wedge n) = (x \cap y) \cap s.$$

This implies  $x \cap y \in s$ , and hence

$$x \bar{\wedge} y = (x \cap y) \cup (x \cap 0) \cup (y \cap 0) = x \cap y;$$

in other words,  $(S_n)_0 = S$ .

Finally, suppose that  $n$  is neutral in  $S$ . Since  $0$  is neutral in  $S_n$ ,

$$\begin{aligned} ((x \cap 0) \cup (y \cap 0)) \cap ((y \cap 0) \cup (z \cap 0)) &= (x \bar{\wedge} y) \cap (y \bar{\wedge} z) \cap 0 = \\ &= (x \cap y) \cap (y \cap z) \cap 0 = [(x \cap y) \cap (y \cap z)] \cap n = \\ &= (x \cap y \cap n) \vee (y \cap z \cap n) = n \wedge j(x, y, z) \end{aligned}$$

as  $n$  is neutral. Also it can be easily shown that  $x \cap y, y \cap z \in$   
 $\in j(x, y, z) = J_0(x, y, z)$ . Therefore

$$[((x \cap 0) \cup (y \cap 0)) \cap ((y \cap 0) \cup (z \cap 0))] \cup (x \cap y) \cup (y \cap z)$$

exists in  $S_n$ ; whence  $0$  is sesquimedial in  $S_n$ . The rest follows by  
 2.6.  $\square$

It should be noted that the above proposition is not true  
 when  $n$  is merely nearly neutral. For example, in Figure 2 which  
 is the isotope of Figure 1,  $0$  is not sesquimedial.

3. Medial nearlattices. Recall that a nearlattice  $S$  is medial if for all  $x, y, z \in S$ ,  $m(x, y, z) = (x \wedge y) \vee (y \wedge z) \vee (z \wedge x)$  exists. A nearlattice  $S$  is said to have the three property if, for any  $x, y, z \in S$ ,  $x \vee y \vee z$  exists whenever  $x \vee y$ ,  $y \vee z$  and  $z \vee x$  exist. Nearlattices with the three property were discussed by Evans in [3], where he referred to them as strong conditional lattices. It is easy to see that a nearlattice  $S$  has the three property if and only if it is medial.

Lemma 3.1. Every element of a medial nearlattice is sesqui-medial.

Proof. Suppose  $S$  is medial and  $n$  is any element of  $S$ . For any  $x, y, z \in S$ ,  $((x \wedge n) \vee (y \wedge n)) \wedge ((y \wedge n) \vee (z \wedge n))$ ,  $x \wedge y \notin m(x, n, y)$  and  $((x \wedge n) \vee (y \wedge n)) \wedge ((y \wedge n) \vee (z \wedge n))$ ,  $y \wedge z \notin m(y, n, z)$ . Thus using the upper bound property and the three property of  $S$ ,  $((x \wedge n) \vee (y \wedge n)) \wedge ((y \wedge n) \vee (z \wedge n)) \vee (x \wedge y) \vee (y \wedge z) = J_n(x, y, z)$  exists in  $S$ .  $\square$

Suppose  $S$  is a medial nearlattice and  $a, b, c \in S$ . If  $a \vee b$ ,  $b \vee c$ ,  $c \vee a$  exists, we define  $m^d(a, b, c) = (a \vee b) \wedge (b \vee c) \wedge (c \vee a)$ . Of course, when  $S$  is distributive,  $m^d(a, b, c) = m(a, b, c)$ . For a fixed element  $n$  of  $S$ , let us introduce a ternary operation  $M_n$ , defined by  $M_n(x, y, z) = m^d(x \wedge n, y \wedge n, z \wedge n) \vee m(x, y, z)$ ;  $x, y, z \in S$ . Notice that  $m^d(x \wedge n, y \wedge n, z \wedge n)$  always exists in  $S$ . But also we have:

Lemma 3.2. In a medial nearlattice  $S$  with  $n \in S$ ,  $M_n(x, y, z)$  always exists for all  $x, y, z \in S$ .

Proof. Notice that  $m^d(x \wedge n, y \wedge n, z \wedge n)$ ,  $x \wedge y \notin m(x, n, y)$ ,  $m^d(x \wedge n, y \wedge n, z \wedge n)$ ,  $y \wedge z \notin m(y, n, z)$  and  $m^d(x \wedge n, y \wedge n, z \wedge n)$ ,  $z \wedge x \notin m(z, n, x)$ . Then by the upper bound property and the three property both  $m^d(x \wedge n, y \wedge n, z \wedge n) \vee (z \wedge x)$  and  $m^d(x \wedge n, y \wedge n, z \wedge n) \vee (x \wedge y) \vee (y \wedge z)$  exist. Thus a second application of the three property yields the existence of  $M_n(x, y, z)$ .  $\square$

Note that if  $n$  is nearly neutral in a nearlattice  $S$ ,  $M_n(x, y, z) = ((x \wedge n) \wedge (y \wedge n) \wedge (z \wedge n) \wedge n) \vee m(x, y, z)$ , and when  $n$  is neutral,  $M_n(x, y, z) \wedge n = (x \wedge n) \wedge (y \wedge n) \wedge (z \wedge n) \wedge n$ . Also if  $S$  is a lattice and  $n$  is neutral,  $M_n(x, y, z) = (m^d(x, y, z) \wedge n) \vee m(x, y, z) = m^d(x, y, z) \wedge \wedge (n \vee m(x, y, z))$ .

Of course  $m(x, y, z)$  and  $M_n(x, y, z)$  are symmetric in  $x, y$  and  $z$ , whereas  $j(x, y, z)$  and  $J_n(x, y, z)$  are not. Thus, the operations

$m$  and  $M_n$  are better behaved and easier to handle than the operations  $j$  and  $J_n$  respectively.

The following proposition is easily verifiable and so is given without proof.

**Proposition 3.3.** For an element  $n$  of a medial nearlattice  $S$ ,  $M_n(x,y,z) = m(x,y,z)$  for all  $x,y,z \in S$  if and only if  $(n)$  is a distributive lattice.

Hence in a distributive medial nearlattice  $S$ ,  $M_n(x,y,z) = m(x,y,z)$  for all  $x,y,z \in S$ .  $\square$

Now we present the following interesting result which extends Theorem 2.6.

**Theorem 3.4.** Suppose  $n$  is a neutral sesquimedial element of a nearlattice  $S$ . Then the following conditions are equivalent.

- (i)  $S$  is medial;
- (ii)  $S_n$  is a medial nearlattice and  $m^{S_n}(x,y,z) = M_n(x,y,z)$  for all  $x,y,z \in S$ .

Moreover, (i) does not necessarily imply (ii) when  $n$  is merely nearly neutral.

**Proof.** (i)  $\Rightarrow$  (ii). Since  $n$  is neutral,

$$M_n(x,y,z) \wedge n = (x \wedge y) \wedge (y \wedge z) \wedge (z \wedge x) \wedge n.$$

By [2, Th. 2.4],

$$M_n(x,y,z) \equiv m(x,y,z)(\Theta_n)$$

and  $x \wedge y \equiv x \wedge y(\Theta_n)$ . Thus,  $(x \wedge y) \wedge (M_n(x,y,z) \equiv x \wedge y(\Theta_n))$ . Similarly,

$$(y \wedge z) \wedge M_n(x,y,z) \equiv y \wedge z(\Theta_n),$$

and

$$(z \wedge x) \wedge M_n(x,y,z) \equiv z \wedge x(\Theta_n).$$

Then using the technique of the proof of (i)  $\Rightarrow$  (ii) in Theorem 2.6, we obtain  $\langle x \wedge y, y \wedge z, z \wedge x \rangle_n = \langle M_n(x,y,z) \rangle_n$ , and (ii) follows from the isomorphism of  $(S_n; \subseteq)$  and  $(P_n(S); \subseteq)$ .

(ii)  $\Rightarrow$  (i). Adjoin a new 0 in  $S$  and form  $(S;0)_n$ . Then by 2.10, 0 is neutral and medial in  $(S;0)_n$ . Thus  $(S;0)_n$  is medial as  $S_n$  is medial. Hence, by (i)  $\Rightarrow$  (ii),  $((S;0)_n)_0$  is medial. But  $((S;0)_n)_0 = (S;0)$  by 2.10, and so  $S$  is medial as required.

For the final assertion consider the lattice of Figure 1, where  $n$  is nearly neutral but not neutral. But its isotope,

given by Figure 2 is not medial.  $\square$

It is well-known by Kolibiar [6] that if  $L$  is a lattice with  $0$  and  $1$  and  $n$  is central in it, then  $L_n$  is also a bounded lattice with  $n$  and  $n'$  as the smallest and the largest elements respectively, where  $x \vee y = m(x, n', y)$  for all  $x, y \in L$ .

An element  $n$  in a lattice  $L$  is called central if it is neutral and complemented in each interval containing it.

We conclude this paper with the following extension of Kolibiar's result.

Proposition 3.5. Suppose  $L$  is a lattice and  $n \in L$  is standard. Then the isotope  $L_n$  is a lattice if and only if  $n$  is central in  $L$ .

Proof. Since  $n$  is standard,  $(L; \varepsilon)$  and  $(P_n(L); \varepsilon)$  are isomorphic by 2.2. Thus,  $L_n$  is a lattice if and only if  $P_n(L)$  is a lattice, i.e. if and only if  $n$  is complemented in each interval containing it. Consequently, the result follows by [2, Th. 3.5].




Figure 1




Figure 2

#### References

- [1] William H. CORNISH and R.C. HICKMAN: Weakly distributive semilattices, Acta Math. Acad. Sci. Hungar. 32(1978), 5-16.
- [2] William H. CORNISH and A.S.A. NOOR: Standard elements in a nearlattice, Bull. Austral. Math. Soc. 26(1982), 185-213.
- [3] E. Evans: Median lattices and convex subalgebras, Colloq. Math. Soc. Janos Bolyai, 29, North Holland Amsterdam, New York 1982, 225-240.
- [4] Robert HICKMAN: Join algebras, Commun. Alg. 8(1980), 1653-1685.
- [5] J. JAKUBÍK and M. KOLIBIAR: On some properties of a pair

- of lattices (Russian), Czechoslovak Math. J. 4(1954), 1-27.
- [6] Milan KOLIBIAR: A ternary operation in lattices (Russian), Czechoslovak Math. J. 6(1956), 318-329.
- [7] A.S.A. NOOR: Isotopes of nearlattices, Ganit. J. Bangladesh Math. Soc. 1(1)(1981), 63-72.
- [8] M. SHOLANDER: Medians, lattices and trees, Proc. Amer. Math. Soc. 5(1954), 808-812.

Department of Mathematics  
Rajshahi University  
Rajshahi  
Bangladesh

School of Mathematical Sciences  
Flinders University  
Bedford Park  
South Australia 5042  
Australia

(Oblatum 19.11. 1986)