

## Werk

**Titel:** A correction to my paper:"Some topological extensions of plane geometry"

**Autor:** Bell, Harold

**Jahr:** 1976

**PURL:** [https://resolver.sub.uni-goettingen.de/purl?320387429\\_0010|log18](https://resolver.sub.uni-goettingen.de/purl?320387429_0010|log18)

## Kontakt/Contact

[Digizeitschriften e.V.](#)  
SUB Göttingen  
Platz der Göttinger Sieben 1  
37073 Göttingen

✉ [info@digizeitschriften.de](mailto:info@digizeitschriften.de)

**A CORRECTION TO MY PAPER : "SOME TOPOLOGICAL EXTENSIONS OF  
PLANE GEOMETRY "**

by

**Harold BELL**

**THEOREM 4.4** and its proof, in my paper : *Some topological extensions of plane geometry*, Rev. Colombiana Mat., 1975, IX, 125 - 153 , should appear as follows : Let  $B$  be a compact set for which the interior of  $T_K(B)$  contains  $A$ . Let  $\mathcal{J} = \{ J : J \in \mathcal{J}_A(e) \text{ for some } e \in B \}$ , and let  $M = \cup \{ J : J \in \mathcal{J} \} \cup A$ . Then the boundary of  $T_K(M)$  is a simple closed curve provided that  $e(K) \notin \bar{M}$ .

*Proof:* Let  $D_1, D_2, D_3, \dots$  be a sequence of simple closed curves such that  $M \subset T_K(D_{n+1}) \subset T_K(D_n)$  for  $n = 1, 2, 3, \dots$ , and each component of  $D_n - M$  has diameter  $< 1/n$ . Since  $A$  is compact there is a  $d > 0$  such that

$$\{ z : d(z, A) < d \} \subset T_K(B) \text{ and } d(e(K), A) > d.$$

If  $e \in E(A)$  and  $d(e, A) < d/3$  then there is an  $f \in L_e \cap B$ . It follows that  $\cup \mathcal{J}_e(A) \subset T_K(I \cup M)$  for some  $I \in \mathcal{J}_f(A)$ . It follows that for each  $\varepsilon > 0$  there is a positive integer  $N$  such that if  $m \geq N$  and  $C$  is a component of  $D_m - M$  then the diameter of  $(T_K(C \cup M) - T_K(M))$  is less than  $\varepsilon$ . A standard uniform convergence argument together with the fact that  $T_K(M)$  has no cut points then yields that the boundary of  $T_K(M)$  is a simple closed curve.

